



CIGNA MEDICAL COVERAGE POLICY

The following Coverage Policy applies to all health benefit plans administered by CIGNA Companies including plans formerly administered by Great-West Healthcare, which is now a part of CIGNA

Subject Percutaneous Vertebroplasty, Kyphoplasty, and Sacroplasty

Effective Date 4/15/2011
Next Review Date 4/15/2012
Coverage Policy Number 0040

Table of Contents

Coverage Policy	1
General Background	2
Coding/Billing Information	11
References	13
Policy History	17

Hyperlink to Related Coverage Policies

- Intervertebral Disc (IVD) Prostheses
- Mechanical Devices for the Treatment of Back Pain
- Minimally Invasive Treatment of Back Pain

INSTRUCTIONS FOR USE

Coverage Policies are intended to provide guidance in interpreting certain **standard** CIGNA HealthCare benefit plans. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement (GSA), Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document **always supercedes** the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. Proprietary information of CIGNA. Copyright ©2011 CIGNA

Coverage Policy

CIGNA covers percutaneous kyphoplasty as medically necessary when standard medical therapy has failed to alleviate symptoms and ANY of the following criteria is met:

- osteoporotic, osteolytic, osteonecrotic (i.e., Kummell disease), or steroid-induced vertebral compression fracture with persistent, debilitating pain unresponsive to at least six weeks of conservative medical management
- severe back pain secondary to destruction of vertebral body due to osteolytic vertebral metastasis or multiple myeloma
- painful and/or aggressive hemangioma or eosinophilic granuloma of the spine

CIGNA covers percutaneous vertebroplasty as medically necessary when standard medical therapy has failed to alleviate symptoms and EITHER of the following criteria is met:

- severe back pain secondary to destruction of vertebral body due to osteolytic vertebral metastasis or multiple myeloma
- painful and/or aggressive hemangioma or eosinophilic granuloma of the spine

CIGNA does not cover percutaneous vertebroplasty or kyphoplasty for any other indication because each is considered experimental, investigational, or unproven.

CIGNA does not cover percutaneous sacroplasty for any indication because it is considered experimental, investigational, or unproven.

General Background

Percutaneous vertebroplasty is a minimally invasive procedure in which bone cement (e.g., polymethylmethacrylate [PMMA]) is injected into a diseased vertebral body under fluoroscopic and/or computed tomography (CT) guidance. Percutaneous kyphoplasty, also referred to as balloon-assisted vertebroplasty, balloon kyphoplasty, or percutaneous vertebral augmentation, was developed as a modification of vertebroplasty. It includes expansion of the vertebra with an inflatable balloon tamp prior to the injection of bone cement. Both procedures have been used to treat osteoporotic vertebral compression fractures and vertebral fractures resulting from osteolytic destruction secondary to malignancy. Sacroplasty, a more recent variation of vertebroplasty, involves injection of PMMA into the sacrum for the treatment of sacral insufficiency fractures. These procedures have been proposed as alternatives to medical management to alleviate pain, provide spine stability and prevent further vertebral collapse in patients with vertebral compression fractures. Conservative medical management of osteoporotic vertebral fractures may include analgesics, activity modification, bracing, and physical therapy, and medications including calcitonin, strontium ranelate, or ibadronate may be provided in an attempt to prevent future fractures. For patients with osteolytic destruction secondary to malignancy, these procedures have been proposed as alternatives to medical management, localized radiation therapy, and traditional surgical stabilization.

Vertebroplasty and kyphoplasty are contraindicated in burst fractures, which result from extreme force applied straight down on the vertebrae, and involve compression of both the anterior and middle columns. Burst fractures can be unstable if the posterior column has sustained injury, and may result in spinal cord injury. Additional contraindications include pedicle fractures, spinal canal or neural foramen compromise, cortical disruption, infection, myelopathy, coagulopathy, allergy to device or material, radiculopathy symptoms, pregnancy, high energy trauma, severe cardiopulmonary deficiencies, active osteomyelitis of the target vertebra, asymptomatic vertebral body compression fracture of patient improving with medical therapy, and use as prophylaxis in osteoporotic patients.

U.S. Food and Drug Administration (FDA)

Several bone cements received 510(k) approval in 2004–2005 for use in vertebroplasty and/or kyphoplasty, including KyphX[®] HV-R[™] Bone Cement (Kyphon Inc., Sunnyvale, CA); Symphony[™] VR Radiopaque Bone Cement (Advanced Biomaterial Systems, Inc. Chatham, NJ; and Parrallax[®] Acrylic Resin with TRACERS[®] (ArthroCare Corp., Sunnyvale, CA. Numerous additional manufacturers subsequently received 510(k) FDA approval for bone cement for use in vertebroplasty and kyphoplasty.

Percutaneous Vertebroplasty

Percutaneous vertebroplasty was first reported in France in 1987 as a treatment for complicated vertebral body hemangioma. The procedure has since been investigated as a treatment to provide mechanical support and pain relief for other conditions associated with osteolytic destruction of the spine. Reported uses of percutaneous vertebroplasty include treatment of osteoporotic fractures, vertebral metastasis, vertebral involvement of multiple myeloma, and, less frequently, aggressive vertebral hemangiomas, Langerhans cell histiocytosis, (i.e., eosinophilic granuloma), and vertebral lymphoma. The mechanism of pain relief attributed to vertebroplasty is not well understood. It has been proposed that pain relief is achieved through stabilization of a weakened vertebral body or by thermal damage to intraosseous nerve fibers.

Percutaneous vertebroplasty gained acceptance by some practitioners as a safe and effective method to provide pain relief, increased mobility and improved quality of life for patients with painful osteolytic lesions and osteoporotic compression fractures. This acceptance and subsequent expanded use of the procedure was primarily based on evidence from uncontrolled studies, however. Until recently, the available literature indicated that these procedures could provide effective treatment with few complications in selected patients. It is likely that the placebo effect as well as the natural history of vertebral fractures contributed in some cases to reported improvements, however, and the safety and efficacy of vertebroplasty had not been evaluated in randomized controlled trials. Two randomized sham-controlled trials published in 2009 demonstrated a lack of improved outcomes with vertebroplasty compared to a sham procedure in the treatment of osteoporotic vertebral

compression fractures. These studies constitute the highest quality evidence published to date. In addition, the use of vertebroplasty has been questioned by recently published technology assessments and specialty society guidelines, detailed below. Guidelines from the American Academy of Orthopaedic Surgeons published in 2010, based on a systematic review of studies published in peer reviewed journals, strongly recommend against vertebroplasty for the treatment of osteoporotic spinal compression fractures. Although vertebroplasty is a minimally invasive procedure, it is not without risk of complications. Complications may include localized bleeding, infection, and pain or neurological symptoms following leakage of injected materials. Although rare, injected material may leak into the venous circulation, potentially resulting in pulmonary embolism.

Literature Review: Klazen et al. conducted an open-label randomized trial to determine whether vertebroplasty has additional value compared with optimum pain treatment in patients with acute vertebral fractures. The study used the following inclusion criteria: vertebral compression fractures on spine radiograph, minimum 15% height loss, fracture at T5 or lower, bone edema on magnetic resonance imaging (MRI), back pain for six weeks or less, and a visual analog score (VAS) of five or more. Of 431 patients eligible for randomization, 229 (53%) had spontaneous pain relief during assessment and did not participate. Of the remaining 202 patients with persistent pain, 101 were randomized to vertebroplasty and 101 to conservative treatment. Vertebroplasty resulted in greater pain relief than conservative treatment; the difference in mean VAS score between baseline and one month was -5.2 after vertebroplasty and -2.7 after conservative treatment, and between baseline and one year was -5.7 after vertebroplasty and -3.7 for conservative treatment. Interpretation of the results of this study is difficult; the analysis included a responder analysis as a primary outcome; a clinically significant change in pain was defined a priori as a three point difference on the VAS scale, and although patients couldn't be blinded to group assignments, no attempt was made to blind treatment assignment for those assessing outcome. In addition, 10% of the patients in the conservative treatment crossed over to the vertebroplasty arm, but detail regarding how and when these crossovers occurred is not provided ((WA Health Technology Assessment, 1010; Blue Cross Blue Shield Technology Evaluation Center [TEC], 2011).

Buchbinder et al. (2009) conducted a multicenter, randomized double-blind controlled trial to determine the short-term efficacy and safety of vertebroplasty for alleviating pain and improving physical functioning in patients with osteoporotic vertebral fractures (n=78). Patients with one or two painful osteoporotic vertebral fractures of less than 12 months duration, confirmed as unhealed by magnetic resonance imaging (MRI), were randomly assigned to vertebroplasty (n=38) or a sham procedure (n=40). Outcomes were evaluated at one week and at one, three, and six months. The primary outcome was overall pain at three months. In the vertebroplasty group, the left pedicle of the fracture site was identified, the skin overlying the pedicle was infiltrated with a 25-gauge needle, and the periosteum of the posterior lamina was infiltrated with a 23-gauge needle. An incision was made in the skin, and a 13-gauge needle was placed posterolaterally relative to the eye of the pedicle. Gentle tapping guided the needle through the pedicle into the anterior two thirds of the fractured vertebral body. PMMA was then injected into the vertebral body. Patients in the sham intervention group underwent the same procedure up until the insertion of the 13-gauge needle. To simulate vertebroplasty, the vertebral body was gently tapped, and PMMA was prepared so that its smell permeated the room. Of the 78 enrolled patients, 35 of 38 in the vertebroplasty group and 36 of 40 in the placebo group completed the six month follow-up. Vertebroplasty did not result in a significant advantage in any measured outcome at any time point. There was a significant reduction in overall pain in both groups at each assessment. Similar improvements were seen in both groups for pain at night and at rest, physical functioning, quality of life, and perceived improvement. The authors concluded that no significant benefit of vertebroplasty over a sham procedure was demonstrated after six months of follow-up, and that these findings call into question the use of vertebroplasty in such patients.

Kallmes et al. conducted a multicenter randomized, double-blind controlled trial to evaluate the efficacy of vertebroplasty in the treatment of painful osteoporotic compression fractures (n=131). Patients were randomized to receive vertebroplasty with PMMA (n=68) or a simulated procedure without PMMA (n=63). For all patients, the skin and subcutaneous tissues overlying the pedicle of the target vertebra or vertebrae were infiltrated with lidocaine, and the periosteum of the pedicle was infiltrated with bupivacaine. Patients were then randomly assigned to receive vertebroplasty or the control intervention. In the vertebroplasty group, needles were passed into the central aspect of the target vertebra or vertebrae, and PMMA was infused. During the control intervention, verbal and physical cues (e.g., pressure on the patient's back) were given, and the methacrylate monomer was opened to simulate the odor of PMMA, but the needle was not placed and PMMA was not infused. The primary outcomes were scores on the modified Rolan-Morris Disability Questionnaire (RDQ) (on a scale of 0–23, with higher scores indicating greater disability), and patients' rating of average pain intensity during the preceding 24 hours at one month (on a scale of 0–10, with higher scores indicating more severe

pain). At one month, there was no significant difference between the two groups in the RDQ score ($p=0.49$) or the pain rating ($0=0.19$). Both groups had immediate improvement in disability and pain scores after the intervention. Although the groups did not differ significantly on any secondary outcome measure at one month, there was a trend toward a higher rate of clinically meaningful improvement in pain in the vertebroplasty group ($p=0.06$).

Diamond et al. (2003) conducted a prospective case-control study in which 79 consecutive osteoporotic patients with acute vertebral fractures were treated with percutaneous vertebroplasty ($n=55$) or conservative therapy alone ($n=24$). All patients presented to the emergency department or were inpatients at a hospital in Australia. Patients who met the inclusion criteria were offered vertebroplasty. Patients who declined the procedure and agreed to longitudinal evaluation comprised the control group. The mean follow-up period was 215 days (range 57–399). Twenty-four hours after vertebroplasty there was a 53% reduction in pain scores from 19 to nine, as measured by the VAS, and a 29% improvement in physical functioning from 14 to 18, as measured by Barthel Index. There was no change in VAS and Barthel Index scores for patients in the conservatively treated group at 24 hours. Clinical outcomes at six weeks and 6–12 months were similar in both groups, however. The authors concluded that when compared to conservative therapy, percutaneous vertebroplasty results in prompt pain relief and rapid rehabilitation. The authors also stated that carefully designed and well-executed long-term clinical trials are needed to verify that this procedure is effective and superior to conservative therapy for managing acute osteoporotic vertebral fractures.

Vertebroplasty has been evaluated in several prospective case series for treatment of osteoporotic vertebral fractures. McGraw et al. (2002) reported that 93% of patients achieved significant improvement in back pain as well as improved ambulatory ability at a mean follow-up of 21.5 months. The preoperative mean VAS score was 8.9 ± 1.12 compared to a score of 2.02 ± 1.95 at follow-up. Winking et al. (2004) evaluated treatment-related disability and quality of life using the Oswestry Low Back Pain Disability (OLBPD) Questionnaire. Preoperative OLBPD scores were 3.7 ± 0.2 preoperatively, compared to scores of 1.6 ± 0.1 at six weeks. The median preoperative VAS pain score was seven, compared to median postoperative scores of 1.8 at two days, 2.6 at six weeks, and 2.7 at six months and one year.

The studies described above evaluate vertebroplasty for osteoporotic fractures. Published studies evaluating vertebroplasty for treatment of osteolytic destruction (e.g., metastasis) consist primarily of retrospective case series. Deramond et al. (1998) conducted an early case series evaluating vertebroplasty in 234 patients with pain caused by vertebral hemangioma, osteoporotic collapse or metastasis, or myeloma. The authors reported that percutaneous vertebroplasty was effective in relieving pain and restoring mobility in 92% of patients. These benefits were maintained for patients with nonprogressive conditions, although the period of follow-up was variable, and the details of outcome measures were not reported.

Several additional retrospective case series have evaluated percutaneous vertebroplasty in the treatment of fractures due to malignancy. Alvarez et al. ($n=21$, 2003) reported improvement in VAS score from 9.2 preoperatively to 3.0 at three months, with no major complications. Fourney et al. ($n=34$, 2003) reported improvement in VAS score from eight preoperatively to two at one month. In the study by Chow et al. ($n=15$, 2004) the VAS score improved from 10 with movement and seven at rest to one with movement and zero at rest at 2–12 weeks follow-up. Cement leakage was reported in each of these studies, although most patients had no associated symptoms. These studies demonstrate significant short-term pain relief as measured by VAS. It is difficult to draw conclusions from these studies, however, because of the study design and small number of included patients.

The National Institute for Health and Clinical Excellence (NICE) (United Kingdom) published Interventional Procedure Guidance in 2003, stating that the evidence on the safety and efficacy of percutaneous vertebroplasty appears adequate to support the use of the procedure. The guidance recommends that the procedure only be undertaken when there are arrangements for access to a spinal surgery service, and with prior discussion between a specialist multidisciplinary team that includes a radiologist and spinal surgeon. The guidance further states that the procedure may be used to provide pain relief for patients with severe painful osteoporosis with loss of height and/or compression fractures of the vertebral body, symptomatic vertebral hemangioma, and painful vertebral body tumors (i.e., metastases or myeloma).

Percutaneous Kyphoplasty

Kyphoplasty, also referred to as balloon-assisted vertebroplasty, or balloon kyphoplasty, or percutaneous vertebral augmentation, was introduced in 2001 as a variation of percutaneous vertebroplasty. A specialized bone tamp with an inflatable balloon is used to expand a collapsed vertebral body to approximate its natural height. Mechanical fixation of the vertebral body is performed by injecting PMMA into the expanded cavity using lower pressure, although PMMA leakage outside the vertebral cavity can still occur. Complications of percutaneous kyphoplasty are similar to those seen with vertebroplasty and are relatively rare. Complication rates are highest in patients with malignancy, due to cement leakage from lytic regions in the vertebral bodies. Reported complications are also higher in this population due to poor overall health.

Literature Review: Wardlaw et al. (2009) conducted a multisite randomized controlled trial to assess the efficacy and safety of balloon kyphoplasty in the treatment of painful vertebral fractures (n=300). Fractures were a mean of 5.6 weeks old at randomization in the kyphoplasty group and 6.4 weeks old in the control group. Inclusion criteria consisted of one to three vertebral fractures. At least one fracture was required to have edema assessed by MRI and at least one fracture had to show at least 15% loss of height. Patients were randomized to kyphoplasty treatment (n=149) or to non-surgical care (n=151). The primary outcome was the difference in change from baseline to one month in the short-form (SF)-36 physical component summary score (scale 1–100). One month follow-up was completed by 138 of 149 kyphoplasty patients and 128 of 151 control patients. Mean SF-36 scores improved by 7.2 points at one month in the kyphoplasty group, and by 2.0 points in the non-surgical group (p<0.0001). At 12 months, the difference between kyphoplasty and control had diminished. The authors suggested that improvement in the non-surgical group during the 12 month follow-up was likely due to fracture healing.

Garfin et al. (2006) conducted a prospective multicenter case series to evaluate the safety and effectiveness of kyphoplasty in the treatment of symptomatic vertebral compression fractures. Quality of life was measured using the Standard Form (SF) 36. Mean preoperative scores on five of the eight subscales were significantly diminished, at ≤ 60% of the age and sex-matched normal values. The mean preoperative scores were 6.3 for role physical, 19.0 for bodily pain, 29.9 for physical function and 37.8 for social function. At 24 months follow-up, mean scores improved to 49.7 for role physical, 52.1 for bodily pain, 46.1 for physical function, and 73.5 for social function. The mean index of back function total back disability score improved from 1.4 at baseline to 0.6 at one month, 0.5 at three and twelve months, and 0.4 at 24 months. As judged by the treating physician, extravasation of PMMA outside the vertebral body occurred in 21 of 214 (10%) treated levels. Review of 165 cases by an independent radiologist showed similar findings, with a PMMA extravasation rate of 11%. During the follow-up period, 23 (23%) patients had at least one subsequent painful vertebral compression fracture. Fourteen of these 23 fractures occurred adjacent to a kyphoplasty-treated vertebral body. Of 155 patients enrolled, 55 did not complete the 24 months of follow-up.

Kasperk et al. (2005) conducted a prospective nonrandomized cohort study to evaluate kyphoplasty in the treatment of patients with vertebral fractures and chronic pain of more than 12 months' duration. Patients self-selected to kyphoplasty or standard medical treatment (n=40) or standard medical treatment alone (n=20). All patients were offered kyphoplasty, and those who refused formed the control group. Outcomes were assessed at three and six months. Pain was evaluated using a modified VAS scoring system, and the European Vertebral Osteoporosis Study (EVOS) score was used to evaluate daily activities. Greater improvement in VAS and EVOS scores and fewer pain-related physician visits were reported in the kyphoplasty group than in the control group. The authors concluded that kyphoplasty is a promising addition to current medical treatment in appropriately selected patients but acknowledged that these findings require confirmation by randomized controlled trials.

A prospective nonrandomized comparative study (Grohs, et al., 2005) evaluated the mechanical effects and impact on quality of life over a two year period in patients with vertebral compression fractures treated with kyphoplasty (28 patients/35 kyphoplasties) or vertebroplasty (23 patients/29 vertebroplasties). There were no significant differences between the groups in terms of patient age and age of fractures. In the kyphoplasty group the VAS score was 7.4 before surgery, 3.5 one day after surgery, 3.2 at four months and 2.0 at two years. The OLBD score for the kyphoplasty group was 61% before surgery, decreased to 42% at one year, but increased to 56% at two years. In the vertebroplasty group the VAS score was 7.8 prior to surgery, 3.0 one day after surgery, 5.7 at four months and 5.6 at two years. The OLBD score was 49% before surgery, decreased slightly to 46% at one year, and increased to 52% at two years. The authors concluded that patients with non-recent fractures seem to experience a longer-lasting effect on pain after balloon kyphoplasty, although they experienced a higher

rate of subsequent compression fractures in adjacent levels. Although the decrease in pain persisted at two years, improvement in disability was limited to one year.

Lane et al. (2004) conducted a prospective case series to evaluate kyphoplasty in the treatment of vertebral fractures due to multiple myeloma (n=19). The authors reported meaningful improvement in 16 of 19 patients at three months. OLB scores improved from 49 ± 16.6 to 32.6 ± 13.6 . Improvement in vertebral body height was also reported. The results were compared with those of a cohort of 26 patients treated for osteoporotic compression fractures. There was no difference in OLB improvement between the two groups, but greater anterior vertebral height restoration was seen in the osteoporotic group. The authors concluded that kyphoplasty is a safe treatment for myeloma-related vertebral compression fractures with efficacy in terms of pain relief and functional outcomes that are comparable to results of patients with osteoporosis. The authors also stated that it is unclear how many levels with fractures related to multiple myeloma should have kyphoplasty, or whether vertebral bodies that show ongoing deformity but are painless are candidates for kyphoplasty, and that well-controlled prospective studies are needed to answer these questions.

Several small case series focused on kyphoplasty to treat vertebral fractures due to malignancy. Dudeney et al. (2002) conducted a prospective case series evaluating kyphoplasty in 18 patients (55 vertebral bodies) with vertebral compression fractures resulting from multiple myeloma. At a mean follow-up of 7.4 months, improvements from baseline in SF 36 scores for pain (from 33.2 to 55.4), physical function (from 21.3 to 50.6), and social functioning (from 31.3 to 47.5) were reported. The authors also reported that on average, 34% of height lost at the time of fracture was restored.

Several additional case series evaluated kyphoplasty in the treatment of vertebral fractures reported improvement in pain and functional scores at short-term follow-up ranging from one week to 18 months (Rhyne et al., 2004; Lieberman et al., 2001; Ledlie and Renfro, 2003; Phillips et al., 2003; Gaitanis, et al., 2005; Ledlie and Renfro, 2006).

Revised Interventional Procedure Guidance on balloon kyphoplasty, published by the National Institute for Health and Clinical Excellence (NICE) (UK) in 2006 concluded that current evidence on the safety and efficacy of the procedure appears adequate to support the use of this procedure in the treatment of vertebral compression fractures. The guidance recommends that the procedure be undertaken only with prior discussion by a specialist multidisciplinary team that includes a radiologist and spinal surgeon, and when there are facilities for good imaging and access to a spinal surgery service.

Percutaneous Sacroplasty

Percutaneous sacroplasty, a variation of vertebroplasty, is an evolving technique that has been proposed for the treatment of sacral insufficiency fractures. Two bone trocars are placed between the sacral foramen and sacroiliac joint on the side of the fractured ala, at a 45 degree angle toward the sacroiliac joint. The needles are then inserted to approximately the midpoint of the sacrum, and PMMA is injected through each trocar.

Literature Review:

Bayley et al. (2009) performed a review of the literature to identify the various techniques used for surgical treatment of sacral insufficiency fractures and to evaluate their outcomes. The techniques described included sacroplasty with or without augmented iliosacral screws. No level I, II, or III evidence was available, and only five articles provided follow-up of one year or more. At total of 108 patients were included in the analysis. The authors concluded that results of cement augmentation techniques such as sacroplasty are promising, with immediate pain relief and maintenance of benefit in the medium-to long-term, but questions remain. The optimal technique and long term outcomes of this procedure need further analysis. The authors stated that future prospective clinical studies with an independent observer to analyze the long-term success rate and complications of this procedure are warranted.

Frey et al. published two case series to assess sacroplasty for the treatment of sacral insufficiency fractures. A study published in 2008 evaluated outcomes and complication rates in 52 patients with incapacitating lumbar and/or gluteal pain, with failure of or intolerance to conservative measures. The mean VAS was 8.1 at baseline, 3.6 thirty minutes following the procedure, 2.5 at two weeks, 2.1 at four weeks, 1.7 at 12 weeks, 1.4 at 24 weeks, and 0.8 at 52 weeks. Improvement was statistically significant. The case series published in 2007 evaluated the safety and efficacy of sacroplasty in 37 patients with sacral insufficiency fractures. VAS scores were monitored periodically for one year, and analgesic usage and patient satisfaction were assessed at the

final follow-up. The mean VAS score was 7.7 at baseline, 3.2 within 30 minutes, 2.1 at two weeks, 1.7 at four weeks, 1.3 at 12 weeks, 1.0 at 24 weeks, and 0.7 at 52 weeks post-procedure. At baseline, 20 patients were using narcotic analgesics compared to 12 patients at last follow-up. The procedure was terminated in one patient who developed radicular pain prior to the injection of PMMA that persisted following the procedure. The authors noted in both studies that the natural history of osteoporotic sacral insufficiency fractures is gradual improvement starting within one to two weeks of treatment initiation, but considered it unlikely that regression toward the mean accounted for the rapid pain reduction seen in this study. The authors also acknowledged in both studies that, because of the lack of a control group, a placebo effect could not be excluded.

Systematic Reviews

Eck et al. (2008) conducted a meta-analysis of the literature to evaluate pain relief and risk of complications associated with vertebroplasty vs. kyphoplasty. Studies included prospective randomized and non-randomized trials, retrospective reviews, and case reports. Studies that provided specific VAS scores were included, and studies and case reports that provided information on complications were also included. A total of 168 studies met the inclusion criteria. Mean pre-and postoperative VAS scores for vertebroplasty were 8.36 and 2.68, respectively, with a mean change of 5.68 ($p < .001$). Mean pre-and postoperative VAS scores for kyphoplasty were 8.06 and 3.46, respectively, with a mean change of 4.60 ($p < .001$). Greater improvement was seen with vertebroplasty vs. kyphoplasty ($p < .001$). The risk of new fractures was 17.9% with vertebroplasty compared to 14.1% with kyphoplasty ($p < .01$). The risk of cement leak was lower with kyphoplasty than vertebroplasty, at 7.0% vs. 19.7%, respectively ($p < .001$). The majority of cement leaks were asymptomatic, incidental findings.

Gill et al. (2007) conducted a systematic review and meta-analysis comparing pain reduction following vertebroplasty and kyphoplasty performed for osteoporotic vertebral compression fractures. Fourteen vertebroplasty and seven kyphoplasty prospective and retrospective trials qualified for inclusion, representing 1046 vertebroplasty patients and 263 kyphoplasty patients. Both procedures resulted in a reduction of greater than five points in the VAS in the immediate postoperative period ($p < 0.00001$). The difference in pain relief was not significant between the two procedures. At final follow-up, the VAS for both vertebroplasty and kyphoplasty was improved, but this improvement was not considered significant when compared to the preoperative VAS ($p = 0.25$, $p = 0.38$, respectively). The authors stated that randomized controlled trials are needed to determine which procedure is most effective for vertebral compression fractures.

Bouza et al. (2006) conducted a systematic review of the literature and meta-analysis of clinical studies evaluating the efficacy and safety of balloon kyphoplasty in the treatment of vertebral compression fractures. A total of 26 studies met the inclusion criteria. The results of the clinical series indicate significant improvements in pain intensity, vertebral height, sagittal alignment, functional impairment and quality of life, although studies displayed considerable methodological limitations. The authors also reported that, compared with vertebroplasty, kyphoplasty reduced the loss of height and kyphotic deformity with a lower leakage rate. Leakage affected 7% of patients; complications were seen in 2% of patients; and new vertebral fractures were seen in 16% of patients. The authors of this systematic review concluded that available evidence suggests that kyphoplasty can be effective and safe for the treatment of vertebral compression fractures, but existing studies demonstrate substantial methodological limitations and relatively short follow-up periods. The authors stated, "Better clinical research is required to determine the capacity of balloon kyphoplasty to avoid the functional and physiological sequelae of vertebral compression fractures and to define the true role of the technique among the existing therapeutic options."

Hulme et al. (2006) conducted a systematic review of 69 clinical studies to evaluate the safety and efficacy of vertebroplasty and kyphoplasty. The authors reported that some pain relief was obtained in 87% of vertebroplasty patients and 92% of kyphoplasty patients. Vertebral height restoration was possible using kyphoplasty (average 6.6 degrees) and a subset of vertebroplasty patients (average 6.6 degrees). Cement leakage was the most common complication, and was reported in 41% of vertebroplasty procedures and 9% of kyphoplasty procedures. The authors stated that, although considerable cement leakage has been reported, the actual rate of leakage may have been underreported because of reporting bias or the detection method used. Cement leakage was asymptomatic in 96% of vertebroplasty patients and 89% of kyphoplasty patients. The authors stated that, until asymptomatic cement leakage can be disregarded as irrelevant, CT is the best method to monitor leakage. New fractures of adjacent vertebrae occurred following both procedures. The authors stated that it has not been determined whether this is caused by altered loading, increased patient activity or the natural progression of osteoporosis. The authors concluded that it cannot be definitively stated that vertebroplasty and kyphoplasty are safe and effective procedures because of a lack of comparative, blinded,

randomized clinical trials. The authors further stated that the adoption of standardized reporting and methodology and an increase in methodological quality would enhance comparability, validity and applicability of the studies.

Taylor et al. (2006) conducted a comparative systematic review of the efficacy and safety of balloon kyphoplasty (four nonrandomized comparative studies and 13 case series) and vertebroplasty (two nonrandomized comparative studies and 57 case series) for vertebral compression fractures, and updated the review in 2007 to include four additional comparative studies and 21 case series. Five studies compared kyphoplasty to vertebroplasty and three to conventional medical care for 481 fractures in 313 patients. The case series included 2047 patients treated with kyphoplasty on 3301 vertebral levels. The median duration of follow-up ranged from hours up to 37 months post-procedure. The limited level of reporting of methods hindered evaluation of methodological quality. The studies that directly compared kyphoplasty to medical care demonstrated significant pain reduction with kyphoplasty, as measured by VAS at 3, 6, 12 and 36 months, and this reduction in pain was greater with kyphoplasty than with medical care alone. Five studies compared kyphoplasty to vertebroplasty. Both procedures reduced VAS pain and improved Oswestry disability index scores for up to 24 months post-procedure, with no significant difference between procedures. All case series in the systematic review reported reduction in pain after kyphoplasty. Studies that evaluated vertebral height and kyphotic angle correction reported improvements in favor of kyphoplasty, with an average improvement in vertebral height of 21% and reduction of 6.3 degrees in kyphotic angle. Safety outcomes of kyphoplasty were combined for comparative studies and case series. A total of 28 studies included details of cement leakages, and eight of these reported whether the leaks were symptomatic or not. A total of 189 (9%) cement leakages were reported in 2239 vertebrae. One leak was reported to be symptomatic, resulting in radiculopathy. A total of 171 new or incident fractures were reported in 1151 patients/16 studies, with 64% of these occurring in adjacent vertebrae. Two of the studies reported fewer subsequent fractures in patients treated with kyphoplasty, however, compared to those who received conventional medical care. The authors concluded that kyphoplasty is more effective than medical management of osteoporotic vertebral compression fractures and is at least as effective as vertebroplasty.

Technology Assessments

Blue Cross Blue Shield Association Technology Evaluation Center (TEC): A BCBS TEC Assessment (in press, Feb, 2011) evaluated the available evidence to determine whether either percutaneous vertebroplasty or kyphoplasty improved the net health outcome for individuals with painful vertebral fractures caused by osteoporosis. The assessment updated the 2010 assessment, incorporating additional studies, including the double-blind, randomized sham control trials by Kallmes (2010) and Buchbinder (2010), and the open label randomized trial by Klazen (2010), all described above. The assessment concluded that the two randomized double-blind controlled trials provide evidence that vertebroplasty may not improve health outcomes when compared to a sham procedure. Because both trials were underpowered, these results should be interpreted as uncertainty rather than a lack of effect. The trial by Klazen investigated vertebroplasty in patients with acute fracture, unlike the participants enrolled in the placebo-controlled trials, and due to methodological issues, interpretation of these data is unclear. The assessment also concluded that the evidence is insufficient to determine if vertebroplasty improves the net health outcome or is as beneficial as any established alternatives. For these reasons, vertebroplasty for vertebral fractures from osteoporosis does not meet the TEC criteria.

The TEC assessment conclusions regarding kyphoplasty remain unchanged from the previous assessment. It is not possible to quantify the benefit of the procedure, or to determine if kyphoplasty improves the net health outcome or is as beneficial as any established alternative.

Washington State Health Care Authority Health Technology Assessment: A Washington State Health Technology Assessment, Vertebroplasty, Kyphoplasty and Sacroplasty, was published in November, 2010, based on a structured, systematic search of the peer reviewed literature. In summarizing the purpose of the review, the authors noted that these procedures are less invasive than other spinal surgical procedures, but more invasive than conservative medical therapy, and although non-randomized studies have reported improvements in pain and functioning, significant questions remain about the safety, efficacy, effectiveness, and cost-effectiveness of these procedures.

The assessment provided the following conclusions:

Efficacy

- Vertebroplasty
 - Pain relief: Uncertain whether vertebroplasty is effective for pain relief.
 - Function and quality of life: Uncertain whether vertebroplasty improves patient functioning and quality of life
- Kyphoplasty
 - Pain relief: Uncertain whether kyphoplasty is effective for pain relief.
 - Function and quality of life: Uncertain whether kyphoplasty improves patient functioning and quality of life
- Vertebroplasty compared with kyphoplasty
 - Pain relief: A single poor-quality RCT found equal improvement in back pain scores over six months. The strength of evidence was noted to be very low.
 - Function and quality of life: No evidence of efficacy for these outcomes.
- Sacroplasty: No evidence of efficacy since the only available evidence consists of case series.

Effectiveness

- Vertebroplasty
 - Pain relief: Uncertain whether vertebroplasty is more effective than conservative medical treatment in reducing pain. Four nonrandomized studies with follow-up of up to a year found that vertebroplasty was more effective than conservative medical treatment up to approximately six months. Pain levels were comparable at one year in both groups. The strength of evidence was noted to be very low.
 - Function and quality of life: A similar pattern was seen in improvements in these four studies in functioning and quality of life, with superior effectiveness in the first 3-6 months followed by equivalent levels of functioning at one year. The strength of evidence was noted to be very low.
- Kyphoplasty:
 - Pain relief: In two non-randomized studies, kyphoplasty reduced pain more than conservative medical treatment for periods up to three years.
 - Function and quality of life: In these two studies, kyphoplasty improved a limited set of functional outcomes more than conservative medical treatment.
- Vertebroplasty compared with kyphoplasty
 - Pain relief: In 8 of 10 non-randomized studies, vertebroplasty and kyphoplasty led to comparable pain reduction up to 2 years.
 - Function and quality of life: In 4 of 5 non-randomized studies, vertebroplasty and kyphoplasty patients demonstrated comparable improvements in ODI up to 2 years.
- Sacroplasty: Unable to draw conclusions due to very limited data.

Regarding safety, the authors stated that, while it appears that the rate of serious complications with associated symptoms are low for vertebroplasty and kyphoplasty, studies with long-term follow up for greater than five years are few, and comparative studies, especially randomized controlled trials, may have too few patients to detect more rare but serious outcomes.

Professional Societies/Organizations: The American Academy of Orthopaedic Surgeons (AAOS) published a clinical practice guideline and evidence report on the treatment of symptomatic osteoporotic spinal compression fractures in 2010. The guideline was based on a systematic review of studies published in English in peer reviewed journals in or after 1966. Additional study requirements included the following: enrollment of ten or more patients per group, results presented quantitatively; enrolled patients 18 years of age or older; not an in vitro, biomechanical or cadaver study, results for patients with osteogenesis imperfecta or solid metastatic tumors of the spine excluded or reported separately; at least 50% patient follow-up (in studies with > 50% but

< 80% follow-up, the study quality was downgraded). Results reported as post-hoc subgroup analyses were excluded. The guideline includes the following recommendations regarding vertebroplasty and kyphoplasty::

- We recommend against vertebroplasty for patients who present with an osteoporotic spinal compression fracture on imaging with correlating clinical signs and symptoms and who are neurologically intact. (Strength of recommendation: strong)
- Kyphoplasty is an option for patients who present with an osteoporotic spinal compression fracture on imaging with correlating clinical signs and symptoms and who are neurologically intact. (Strength of recommendation: weak)

Additional AAOS recommendations regarding treatment of osteoporotic compression fractures include:

- We suggest patients who present with an osteoporotic spinal compression fracture on imaging with correlating clinical signs and symptoms suggesting an acute injury (0-5 days after identifiable event or onset of symptoms) and who are neurologically intact be treated with calcitonin for 4 weeks. (Strength of recommendation: moderate)
- Ibandronate and strontium ranelate are options to prevent additional symptomatic fractures in patients who present with an osteoporotic spinal compression fracture on imaging with correlating clinical signs and symptoms. (Strength of recommendation: weak)
- It is an option to treat patients who present with an osteoporotic spinal compression fracture at L3 or L4 on imaging with correlating clinical signs and symptoms suggesting an acute injury and who are neurologically intact with an L2 nerve root block. (Strength of recommendation: weak)

The authors were unable to recommend for or against the following treatments (Strength of recommendation: inconclusive):

- Bed rest, complementary and alternative medicine, or opioids/analgesics for patients who present with an osteoporotic spinal compression fracture on imaging with correlating clinical signs and symptoms and who are neurologically intact. (Strength of recommendation: inconclusive)
- Treatment with a brace for patients who present with an osteoporotic spinal compression fracture on imaging with correlating clinical signs and symptoms and who are neurologically intact. (Strength of recommendation: inconclusive)
- A supervised or unsupervised exercise program for patients who present with an osteoporotic spinal compression fracture on imaging with correlating clinical signs and symptoms and who are neurologically intact. (Strength of recommendation: inconclusive)
- Electrical stimulation for patients who present with an osteoporotic spinal compression fracture on imaging with correlating clinical signs and symptoms and who are neurologically intact. (Strength of recommendation: inconclusive)

A practice guideline for the performance of vertebroplasty developed in collaboratively by the American College of Radiology, American Society of Neuroradiology, Society of Neurointerventional Surgery, American Society of Spine Radiology, and the Society of Interventional Radiology, was revised in 2009. The guideline states that the major indication for vertebroplasty is the treatment of symptomatic osteoporotic or neoplastic vertebral body compression fracture(s) refractory to medical therapy. Failure of medical therapy is defined as minimal or no pain relief with the administration of prescribed analgesics, or adequate pain relief with narcotic dosages that produce undesirable side effects. The guideline includes the following indications for vertebroplasty:

- Painful osteoporotic or neoplastic vertebral compression fracture(s) refractory to medical therapy
- Symptomatic vertebral body microfracture (as documented by magnetic resonance imaging [MRI] or nuclear imaging, and/or lytic lesion seen on CT) without obvious loss of vertebral body height

A 2007 position statement on percutaneous vertebral augmentation developed by the American Society of Interventional and Therapeutic Neuroradiology, Society of Interventional Radiology, American Association of Neurological Surgeons/Congress of Neurological Surgeons, and the American Society of Spine Radiology, states that percutaneous vertebral augmentation with vertebroplasty and kyphoplasty is a safe, efficacious, and durable procedure in appropriate patients with symptomatic osteoporotic and neoplastic fractures when

performed in a manner in accordance with published standards. These procedures are offered only when traditional medical therapy has not provided pain relief or pain is substantially altering the patient's lifestyle.

There are no published specialty society statements or guidelines that address sacroplasty.

Summary

Percutaneous vertebroplasty and kyphoplasty gained acceptance by some practitioners for the treatment of painful osteolytic lesions and osteoporotic compression fractures based on evidence from prospective and retrospective case series. The safety and efficacy of vertebroplasty had not been evaluated in randomized controlled trials, however. Two randomized controlled trials published in 2009 demonstrated a lack of improved outcomes with vertebroplasty compared to a-sham procedure in patients with osteoporotic vertebral fractures. These studies constitute the highest quality evidence published to date. Guidelines from the American Academy of Orthopaedic Surgeons published in 2010, based on a systematic review of studies published in peer reviewed journals, strongly recommend against vertebroplasty for the treatment of osteoporotic spinal compression fractures. Therefore, based on this newly published evidence and specialty society opinion, as well as recently published technology assessments, vertebroplasty has not been demonstrated to result in improved outcomes compared to conservative medical treatment in patients with osteoporotic vertebral compression fractures. Vertebroplasty may be a reasonable treatment option, however, for selected patients with severe back pain secondary to destruction of the vertebral body due to osteolytic vertebral metastasis or multiple myeloma, or for painful and/or aggressive hemangioma or eosinophilic granuloma of the spine, when conservative medical management has failed to alleviate symptoms.

Kyphoplasty, also referred to as percutaneous vertebral augmentation, is a modification of vertebroplasty which includes expansion of the vertebra with an inflatable balloon tamp prior to the injection of bone cement. Although the published medical literature evaluating kyphoplasty is not robust, kyphoplasty has been shown in several non-randomized studies to reduce pain more than conservative medical treatment for periods of up to three years, and improved specific functional outcomes more than conservative medical treatment. Additional well designed clinical trials with long term follow-up are needed to define patient selection criteria and to determine the long-term safety and efficacy of this procedure. Kyphoplasty should therefore be reserved for carefully selected patients in whom conservative medical management has failed to alleviate symptoms.

Sacroplasty, a variation of vertebroplasty, is an evolving technique that has been proposed for the treatment of sacral insufficiency fractures. There is insufficient evidence in the published medical literature to demonstrate the safety, efficacy, and long-term outcomes of this procedure.

Coding/Billing Information

Note: This list of codes may not be all-inclusive.

Percutaneous Kyphoplasty

Covered when medically necessary:

CPT®* Codes	Description
22523	Percutaneous vertebral augmentation, including cavity creation (fracture reduction and bone biopsy included when performed) using mechanical device, one vertebral body, unilateral or bilateral cannulation (e.g., kyphoplasty); thoracic
22524	Percutaneous vertebral augmentation, including cavity creation (fracture reduction and bone biopsy included when performed) using mechanical device, one vertebral body, unilateral or bilateral cannulation (eg, kyphoplasty); lumbar
22525	Percutaneous vertebral augmentation, including cavity creation (fracture reduction and bone biopsy included when performed) using mechanical device, one vertebral body, unilateral or bilateral cannulation (eg, kyphoplasty); each additional thoracic or lumbar vertebral body (List separately in addition to code for primary procedure)

ICD-9-CM Diagnosis Codes	Description
170.2	Malignant neoplasm of bone and articular cartilage; vertebral column excluding sacrum and coccyx
198.5	Secondary malignant neoplasm of bone and bone marrow
203.00	Multiple myeloma without mention of remission
203.01	Multiple myeloma in remission
203.02	Multiple myeloma in relapse
228.09	Hemangioma of other sites
277.89	Other specified disorders of metabolism
721.78	Traumatic spondylopathy
733.13	Pathologic fracture of vertebrae
805.2	Fracture of vertebral column without mention of spinal cord injury, dorsal (thoracic) closed
805.4	Fracture of vertebral column without mention of spinal cord injury, lumbar closed

Experimental/Investigational/Unproven/Not Covered:

ICD-9-CM Diagnosis Codes	Description
	All other codes

Percutaneous Vertebroplasty

Covered when medically necessary:

CPT [®] * Codes	Description
22520	Percutaneous vertebroplasty, one vertebral body, unilateral or bilateral injection, thoracic
22521	Percutaneous vertebroplasty, one vertebral body, unilateral or bilateral injection, lumbar
22522	Percutaneous vertebroplasty, one vertebral body, unilateral or bilateral injection; each additional thoracic or lumbar vertebral body (List separately in addition to code for primary procedure)

HCPCS Codes	Description
S2360	Percutaneous vertebroplasty, one vertebral body, unilateral or bilateral injection; cervical
S2361	Each additional cervical vertebral body

ICD-9-CM Diagnosis Codes	Description
170.2	Malignant neoplasm of bone and articular cartilage; vertebral column excluding sacrum and coccyx
198.5	Secondary malignant neoplasm of bone and bone marrow
203.00	Multiple myeloma without mention of remission
203.01	Multiple myeloma in remission
203.02	Multiple myeloma in relapse
228.09	Hemangioma of other sites

277.89	Other specified disorders of metabolism
733.13	Pathologic fracture of vertebrae
805.00-805.08	Fracture of vertebral column without mention of spinal cord injury; cervical, closed
805.2	Closed fracture of dorsal (thoracic) vertebra without mention of spinal cord injury
805.4	Closed fracture of lumbar vertebra without mention of spinal cord injury

Experimental/Investigational/Unproven/Not Covered:

ICD-9-CM Diagnosis Codes	Description
721.78	Traumatic spondylopathy
	All other codes

Percutaneous Sacroplasty

Experimental/Investigational/Unproven/Not Covered:

CPT* Codes	Description
0200T	Percutaneous sacral augmentation (sacroplasty), unilateral injection(s), including the use of a balloon or mechanical device (if utilized), 1 or more needles
0201T	Percutaneous sacral augmentation (sacroplasty), bilateral injections, including the use of a balloon or mechanical device (if utilized), 2 or more needles

ICD-9-CM Diagnosis Codes	Description
	All codes

*Current Procedural Terminology (CPT®) © 2010 American Medical Association: Chicago, IL.

References

1. Bayley E, Srinivas S, Boszczyk BM. Clinical outcomes of sacroplasty in sacral insufficiency fractures: a review of the literature. *Eur Spine J.* 2009 Sep;18(9):1266-71. Epub 2009 Jun 6.
2. Berlemann U, Franz T, Orlor R, Heini PF. Kyphoplasty for treatment of osteoporotic vertebral fractures: a prospective non-randomized study. *Eur Spine J.* 2004 Oct;13(6):496-501.
3. BlueCross BlueShield Association (BCBSA) Technology Evaluation Center (TEC). TEC in press. Percutaneous vertebroplasty or kyphoplasty for vertebral fractures caused by osteoporosis TEC Assessment Program. Chicago IL: BCBSA; 2011 Feb. Accessed Mar 5, 2011. Available at URL address: <http://www.bcbs.com/blueresources/tec/press/>
4. BlueCross BlueShield Association (BCBSA) Technology Evaluation Center (TEC). Percutaneous vertebroplasty or kyphoplasty for vertebral fractures caused by osteoporosis. TEC Assessment Program. Vol. 24, No. 7. Chicago IL: BCBSA; 2010 Apr. Accessed Mar 8, 2011. Available at URL address: <http://www.bcbs.com/blueresources/tec/tec-assesments-by-topic.html?topics=orthopedicspain-management>
5. Bouza C, Lopez T, Magro A, Navalpotro L, Amate JM. Efficacy and safety of balloon kyphoplasty in the treatment of vertebral compression fractures: a systematic review. *Eur Spine J.* 2006 Jan 21:1-18 [Epub ahead of print]

6. Buchbinder R, Osborne RH, Ebeling PR, Wark JD, Mitchell P, Wriedt C, et al. randomized trial of vertebroplasty for painful osteoporotic vertebral fractures. *N Engl J Med*. 2009 Aug 6;361(6):557-68.
7. Chow E, Holden L, Danjoux C, Yee A, Vidmar M, Connolly R, et al. Successful salvage using percutaneous vertebroplasty in cancer patients with painful spinal metastases or osteoporotic compression fractures. *Radiother Oncol*. 2004 Mar;70(3):265-7.
8. Coumans JVCE, Reinhardt MK, Lieberman IH. Kyphoplasty for vertebral compression fractures: 1-year clinical outcomes from a prospective study. *J Neurosurg*. 2003 Jul;99(1 Suppl):44-50.
9. Crandall D, Slaughter D, Hankins PJ, Moore C, Jerman J. Acute versus chronic vertebral compression fractures treated with kyphoplasty: early results. *Spine J*. 2004 Jul-Aug;4(4):418-24.
10. De Negri P, Tirri T, Paternoster G, Modano P. Treatment of painful osteoporotic or traumatic vertebral compression fractures by percutaneous vertebral augmentation procedures: a nonrandomized comparison between vertebroplasty and kyphoplasty. *Clin J Pain*. 2007 Jun;23(5):425-30.
11. Deramond H, Depriester C, Galibert P, Le Gars, D. Percutaneous vertebroplasty with polymethylmethacrylate. Technique, indications, and results. *Radiol Clin North Am*. 1998 May;36(3):533-46.
12. Deramond H, Saliou G, Aveillan M, Lehmann P, Vallee JN. Respective contributions of vertebroplasty and kyphoplasty to the management of osteoporotic vertebral fractures. *Joint Bone Spine*. 2006 Dec;73(6):610-3. Epub 2006 Oct 11.
13. DeVita VT, Lawrence TS, Rosenberg SA. DeVita, Hellman & Rosenberg's cancer: principles & practice of oncology. Lippincott Williams & Wilkins; 2008.
14. Diamond TH, Champion B, Clark WA. Management of acute osteoporotic vertebral fractures: a nonrandomized trial comparing percutaneous vertebroplasty with conservative therapy. *Am J Med*. 2003 Mar;114(4):257-65.
15. Dudeney S, Lieberman IH, Reinhardt MK, Hussein M. Kyphoplasty in the treatment of osteolytic vertebral compression fractures as a result of multiple myeloma. *J Clin Oncol*. 2002 May 1;20(9):2382-7.
16. Eck JC, Nachtigall D, Humphreys SC, Hodges SD. Comparison of vertebroplasty and balloon kyphoplasty for treatment of vertebral compression fractures: a meta-analysis of the literature. *Spine J*. 2008 May-Jun;8(3):488-97. Epub 2007 May 29.
17. ECRI Institute. Percutaneous vertebroplasty for the treatment of vertebral fractures. Plymouth Meeting (PA): ECRI Institute Health Technology Assessment Information Service; 2008 Dec. Available at URL address: <http://www.ecri.org>
18. ECRI Institute. Percutaneous kyphoplasty for the treatment of vertebral fractures. Plymouth Meeting (PA): ECRI Institute Health Technology Assessment Information Service. 2006 Mar 75 p. (Windows on Medical Technology™, no. 132). Available at URL address: <http://www.ecri.org>
19. Eichholz KM, O'Toole JE, Christie SD, Fessler RG. Vertebroplasty and kyphoplasty. *Neurosurg Clin N Am*. 2006 Oct;17(4):507-18.
20. Fourney DR, Schomer DF, Nader R, Chlan-Fourney J, Suki D, Ahrar K, et al. Percutaneous vertebroplasty and kyphoplasty for painful vertebral body fractures in cancer patients. *J Neurosurg*. 2003 Jan;98(1 Suppl):21-30.
21. Frey ME, Depalma MJ, Cifu DX, Bhagia SM, Carne W, Daitch JS. Percutaneous sacroplasty for osteoporotic sacral insufficiency fractures: a prospective, multicenter, observational pilot study. *Spine J*. 2008 Mar-Apr;8(2):367-73. Epub 2007 Jul 20.

22. Frey ME, DePalma MJ, Cifu DX, Bhagia SM, Daitch JS. Efficacy and safety of percutaneous sacroplasty for painful osteoporotic sacral insufficiency fractures: a prospective, multicenter trial. *Spine (Phila Pa 1976)*. 2007 Jul 1;32(15):1635-40.
23. Gaitanis IN, Hadjipavlou AG, Katonis PG, Tzermiadianos MN, Pasku DS, Patwardhan AG. Balloon kyphoplasty for the treatment of pathological vertebral compressive fractures. *Eur Spine J*. 2005 Apr;14(3):250-60. Epub 2004 Oct 8.
24. Garfin SR, Buckley RA, Ledlie J, Balloon Kyphoplasty Outcomes Group. Balloon kyphoplasty for symptomatic vertebral body compression fractures results in rapid, significant, and sustained improvements in back pain, function, and quality of life for elderly patients. *Spine*. 2006 Sep 1;31(19):2213-20.
25. Grohs JG, Matzner M, Trieb K, Krepler P. Minimal invasive stabilization of osteoporotic vertebral fractures: a prospective nonrandomized comparison of vertebroplasty and balloon kyphoplasty. *J Spinal Disord Tech*. 2005 Jun;18(3):238-42.
26. Hadjipavlou AG, Tzermiadimos MN, Katonis PG, Szpalski M. Percutaneous vertebroplasty and balloon kyphoplasty for the treatment of osteoporotic vertebral compression fractures and osteolytic tumours. *J Bone Joint Surg Br*. 2005 Dec;87(12):1595-604.
27. Hulme PA, Krebs J, Ferguson SJ, Berlemann U. Vertebroplasty and kyphoplasty: a systematic review of 69 clinical studies. *Spine*. 2006 Aug 1;31(17):1983-2001.
28. Jensen ME, McGraw JK, Cardella Jf, Hirsch JA. Position statement on percutaneous vertebral augmentation: a consensus statement developed by the American Society of Interventional and Therapeutic Neuroradiology, Society of Interventional Radiology, American Association of Neurological Surgeons/Congress of Neurological Surgeons, and American Society of Spine Radiology. *Vasc Interv Radiol*. 2009 Jul;20(7 Suppl):S326-31.
29. Kallmes DF, Comstock BA, Heagerty PJ, Turner JA, Wilson DJ, Diamond TH, et al. A randomized trial of vertebroplasty for osteoporotic spinal fractures. *N Engl J Med*. 2009 Aug 6;361(6):569-79.
30. Kasperk C, Grafe IA, Schmitt S, Nöldge G, Weiss C, Da Fonseca K, et al. Three-year outcomes after kyphoplasty in patients with osteoporosis with painful vertebral fractures. *J Vasc Interv Radiol*. 2010 May;21(5):701-9. Epub 2010 Mar 20.
31. Kasperk C, Hillmeier J, Noldge G, Grafe IA, DaFonseca K, Raupp D, et al. Treatment of painful vertebral fractures by kyphoplasty in patients with primary osteoporosis: a prospective nonrandomized controlled study. *J Bone Miner Res*. 2005 Apr;20(4):604-12. Epub 2004 Dec 6.
32. Kaufmann TJ, Jensen ME, Schweickert PA, Marx WF, Kallmes DF. Age of fracture and clinical outcomes of percutaneous vertebroplasty. *AJNR Am J Neuroradiol*. 2001 Nov-Dec;22(10):1860-3.
33. Klazen CA, Lohle PN, de Vries J, Jansen FH, Tielbeek AV, Blonk MC, et al. Vertebroplasty versus conservative treatment in acute osteoporotic vertebral compression fractures (Vertos II): an open-label randomised trial. *Lancet*. 2010 Sep 25;376(9746):1085-92. Epub 2010 Aug 9
34. Lane JM, Hong R, Koob J, Kiechle T, Nievizky R, Pearse R, et al. Kyphoplasty enhances function and structural alignment in multiple myeloma. *Clin Orthop Relat Res*. 2004 Sep;(426):49-53.
35. Lavelle W, Carl A, Lavelle ED, Khaleel MA. Vertebroplasty and kyphoplasty. *Med Clin North Am*. 2007 Mar;91(2):299-314.
36. Ledlie JT, Renfro M. Balloon kyphoplasty: one-year outcomes in vertebral body height restoration, chronic pain, and activity levels. *J Neurosurg*. 2003 Jan;98(1 Suppl):36-42.

37. Ledlie JT, Renfro MB. Kyphoplasty treatment of vertebral fractures: 2-year outcomes show sustained benefits. *Spine*. 2006 Jan 1;31(1):57-64.
38. Legroux-Gerot I, Lormeau, Boutry N, Cotton A, Duquesnoy B, Cortet B. Long-term follow-up of vertebral osteoporotic fractures treated by percutaneous vertebroplasty. *Clin Rheumatol*. 2004 Aug;23(4):310-7. Epub 2004 May 18.
39. Mathis JM, Ortiz AO, Zoarski GH. Vertebroplasty versus kyphoplasty: a comparison and contrast. *AJNR Am J Neuroradiol*. 2004 May;25(5):840-5.
40. McGraw JK, Cardella J, Barr JD, Mathis JM, Sanchez O, Schwartzberg MS, et al.; Society of Interventional Radiology Standards of Practice Committee. Society of Interventional Radiology quality improvement guidelines for percutaneous vertebroplasty. *J Vasc Interv Radiol*. 2003 Sep;14(9 Pt 2):S311-5.
41. McGraw JK, Lippert JA, Minkus KD, Rami PM, Davis TM, Budzik RF. Prospective evaluation of pain relief in 100 patients undergoing percutaneous vertebroplasty: results and follow-up. *J Vasc Interv Radiol*. 2002 Sep;13(9 Pt 1):883-6.
42. McGuire R. AAOS Clinical Practice Guideline: The Treatment of Symptomatic Osteoporotic Spinal Compression Fractures. *J Am Acad Orthop Surg*. 2011 Mar;19(3):183-4. Accessed Mar 3, 2010. Available at URL address: <http://www.aaos.org/Research/guidelines/guide.asp>
43. McKiernan F, Faciszewski T, Jensen R. Reporting height restoration in vertebral compression fractures. *Spine*. 2003 Nov 15;28(22):2517-21; discussion 3.
44. National Institute for Clinical Excellence (NICE). Balloon kyphoplasty for vertebral compression fractures. *Interventional procedure guidance 166*. London, UK: NICE; 2006 Apr. Revised 2008 Jan. Accessed Mar 9 2010. Available at URL address: <http://www.nice.org.uk/Guidance/IPG166/Guidance/pdf/English>
45. National Institute for Clinical Excellence (NICE). Percutaneous vertebroplasty. *Interventional procedure guidance 12*. London, UK: NICE; 2003 Nov. Accessed Mar 9 2010. Available at URL address: <http://www.nice.org.uk/page.aspx?o=85733>
46. Pateder DB, Khanna AJ, Lieberman IH. Vertebroplasty and kyphoplasty for the management of osteoporotic vertebral compression fractures. *Orthop Clin North Am*. 2007 Jul;38(3):409-18; abstract vii.
47. Phillips FM, Ho E, Campbell-Hupp M, McNally T, Todd Wetzel F, Gupta P. Early radiographic and clinical results of balloon kyphoplasty for the treatment of osteoporotic vertebral compression fractures. *Spine*. 2003 Oct 1;28(19):2260-5; discussion 2265-7.
48. Rao RD, Singrakhia MD. Painful osteoporotic vertebral fracture: pathogenesis, evaluation, and roles of vertebroplasty and kyphoplasty in its management. *J Bone Joint Surg Am*. 2003 Oct;85-A(10):2010-22.
49. Richardson P, Hideshima T, Anderson Kc. Multiple myeloma and related disorders. In: *Abeloff: Clinical oncology*. Churchill Livingstone, an imprint of Elsevier; 2004.
50. Rhyne III A, Banit D, Laxwer E, Odum S, Nussman D. Kyphoplasty: report of eighty-two thoracolumbar osteoporotic vertebral fractures. *J Orthop Trauma*. 2004 May-Jun;18(5):294-9.
51. Spivak JM, Johnson MG. Percutaneous treatment of vertebral body pathology. *J Am Acad Orthop Surg*. 2005 Jan-Feb;13(1):6-17.
52. Taylor RS, Fritzell P, Taylor RJ. Balloon kyphoplasty in the management of vertebral compression fractures: an updated systematic review and meta-analysis. *Eur Spine J*. 2007 Feb 3;

53. Taylor RS, Taylor RJ, Fritzell P. Balloon kyphoplasty and vertebroplasty for vertebral compression fractures: a comparative systematic review of efficacy and safety. *Spine*. 2006 Nov 1;31(23):2747-55.
54. Trout AT, Kallmes DF, Kaufmann TJ. New fractures after vertebroplasty: adjacent fractures occur significantly sooner. *AJNR Am J Neuroradiol*. 2006 Jan;27(1):217-23.
55. Wardlaw D, Cummings SR, Van Meirhaeghe J, Bastian L, Tillman JB, Ranstam J, et al. Efficacy and safety of balloon kyphoplasty compared with non-surgical care for vertebral compression fracture (FREE): a randomised controlled trial. *Lancet*. 2009 Mar 21;373(9668):1016-24. Epub 2009 Feb 24.
56. Washington State Health Care Authority. Vertebroplasty, kyphoplasty and Sacroplasty Health Technology Assessment. Olympia WA: Health Technology Assessment Program, 2010 Nov.
57. Winking M, Stahl JP, Oerterl M, Schnettler R, Boker DK. Treatment of pain from osteoporotic vertebral collapse by percutaneous PMMA vertebroplasty. *Acta Neurochir (Wien)*. 2004 May;146(5):469-76. Epub 2004 Apr 8.

Policy History

Pre-Merger Organizations	Last Review Date	Policy Number	Title
CIGNA HealthCare	4/15/2008	0040	Percutaneous Vertebroplasty and Kyphoplasty
Great-West Healthcare	11/30/2007	04.248.03	Vertebroplasty and Kyphoplasty

“CIGNA”, “CIGNA HealthCare” and the “Tree of Life” logo are registered service marks of CIGNA Intellectual Property, Inc., licensed for use by CIGNA Corporation and its operating subsidiaries. All products and services are provided by such operating subsidiaries and not by CIGNA Corporation. Such operating subsidiaries include Connecticut General Life Insurance Company, CIGNA Health and Life Insurance Company, CIGNA Behavioral Health, Inc., CIGNA Health Management, Inc., and HMO or service company subsidiaries of CIGNA Health Corporation and CIGNA Dental Health, Inc. In Arizona, HMO plans are offered by CIGNA HealthCare of Arizona, Inc. In California, HMO plans are offered by CIGNA HealthCare of California, Inc. In Connecticut, HMO plans are offered by CIGNA HealthCare of Connecticut, Inc. In North Carolina, HMO plans are offered by CIGNA HealthCare of North Carolina, Inc. In Virginia, HMO plans are offered by CIGNA HealthCare Mid-Atlantic, Inc. All other medical plans in these states are insured or administered by Connecticut General Life Insurance Company or CIGNA Health and Life Insurance Company