



CIGNA MEDICAL COVERAGE POLICY

The following Coverage Policy applies to all plans administered by CIGNA Companies including plans administered by Great-West Healthcare, which is now a part of CIGNA.

Subject **Ultrasound In Pregnancy
(including 3D and 4D
Ultrasound)**

Effective Date 8/15/2010
Next Review Date 8/15/2011
Coverage Policy Number 0142

Table of Contents

Coverage Policy	1
General Background	1
Coding/Billing Information	4
References	6
Policy History	8

Hyperlink to Related Coverage Policies

Down Syndrome Screening
Fetal Surgery

INSTRUCTIONS FOR USE

Coverage Policies are intended to provide guidance in interpreting certain **standard** CIGNA HealthCare benefit plans as well as benefit plans formerly administered by Great-West Healthcare. Please note, the terms of a participant's particular benefit plan document [Group Service Agreement (GSA), Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a participant's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a participant's benefit plan document **always supercedes** the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable group benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. Proprietary information of CIGNA. Copyright ©2010 CIGNA

Coverage Policy

CIGNA covers one routine two-dimensional (2D) standard obstetrical ultrasound examination during pregnancy. CIGNA covers additional ultrasound examinations as medically necessary when performed for specific medical indications.

CIGNA does not cover an obstetrical ultrasound examination performed solely to determine gender or to provide photographic representation of the fetus, because it is considered not medically necessary for the management of a pregnancy.

CIGNA does not cover either three-dimensional (3D) or four-dimensional (4D) obstetrical ultrasonography because each is considered experimental, investigational or unproven.

General Background

Ultrasound imaging uses high-frequency sound waves to produce dynamic images of organs, tissues or blood-flow inside the body. The procedure involves the use of a transducer, which sends a stream of high-frequency sound waves into the body and detects their echoes as they bounce off internal structures. The sound waves are converted to electrical impulses, which are processed to form an image displayed on a computer monitor. Obstetricians use ultrasounds at a very low power level to check fetal size, location, age and quantity.

Ultrasound is also used in this manner to assess for the presence of some types of birth defects, fetal movement, breathing and heartbeat.

Two-dimensional (2D) ultrasound is considered standard or conventional ultrasound. In conventional 2D scanning the ultrasound image is made up of a series of thin slices and only one slice can be seen at any one time. For three-dimensional (3D) ultrasonography a volume of echoes is taken, which can be stored digitally and shaded to produce life-like pictures of the fetus. It is possible to measure distance, area and volume from volume data with 3D ultrasound. Three-dimensional ultrasound data can be sliced in any orientation, allowing for any diameter or cross-sectional area of the organ to be measured. Four-dimensional ultrasound adds motion to the 3D imaging display. This feature typically involves 3D multiplanar imaging that is acquired at rates that stimulate movement such as heart motion or fetal activity. With 4D ultrasound, the life-like fetal pictures can be seen to move in real time so the activity of the baby inside the womb can be studied.

There is no consensus on the best use of ultrasonography in screening for abnormal pregnancies in low-risk populations. However, many health care providers recommend that one ultrasound examination, usually done between 18 and 20 weeks of pregnancy, be included as a routine part of prenatal care. The use of ultrasonography to assess for potential fetal abnormalities, confirm the site of pregnancy within the uterus, and determine gestational age is considered the standard of care. Also, the use of ultrasound scanning during the first trimester is correlated with reduced post-term labor induction rates as compared to second trimester ultrasound scanning (American College of Obstetricians and Gynecologists [ACOG], 2004a).

ACOG uses the following terms to describe various types of ultrasound examinations performed during the second and third trimesters:

- Standard: includes an evaluation of fetal presentation amniotic fluid volume, cardiac activity, placental position, fetal biometry and an anatomic survey.
- Limited: performed when a specific question requires investigation; appropriate only when the patient has had a prior complete examination.
- Specialized: performed when an anomaly is suspected on the basis of history, biochemical abnormalities or clinical evaluation, or when results from either a limited or standard ultrasound examination are suspicious.

First-Trimester Ultrasound Screening

Obtaining an accurate expected date of delivery (EDD) using ultrasonography early in the pregnancy can reduce the incidence of pregnancies diagnosed as post-term and minimize unnecessary interventions. The premise is that the EDC as calculated by menstrual age is often inaccurate and therefore can be the basis for presumed but incorrect diagnosis of post-term pregnancy. The reported frequency of post-term pregnancy is approximately 7%. Most cases of post-term pregnancy result from a prolongation of gestation. Other cases result from an inability to accurately define EDD. The risk of adverse sequelae may be reduced by making an accurate assessment of gestational age and diagnosis of post-term gestation, as well as recognition and management of risk factors.

Literature Review: Whitworth et al. (2010) conducted a Cochrane review of ultrasound in early pregnancy which included 11 trials (n=37505 women). The analysis found routine scanning to be associated with reduced inductions of labor for post-term pregnancy. Early ultrasound was also found to improve the early detection of multiple pregnancies.

There is sufficient evidence in the published peer-reviewed medical literature (Bennett, et al., 2004; Bukowski, et al., 2003; ECRI, 2003) indicating that the use of ultrasound in early pregnancy (i.e., before 24 weeks gestation) may be of benefit in determining gestational age, detecting multiple pregnancies, and decreasing the rates of induction of labor for post-term pregnancies.

Use of 2D Versus 3D and 4D Ultrasound

The ultimate impact of 3D and 4D ultrasound as new diagnostic imaging technologies is difficult to characterize due to the rapidly changing technological advances in the medical imaging industry. Potential areas of promise include fetal facial anomalies, neural tube defects, and skeletal malformations where 3D ultrasonography may be helpful in diagnosis as an adjunct to, but not a replacement for, 2D ultrasonography (ACOG, 2009). Although 3D ultrasound may provide additional diagnostic information, there is a lack of data demonstrating the impact on

clinical outcomes. Proponents of the use 4D ultrasound suggest that the real-time movements of the fetus obtained improves maternal bonding, however the impact of 4D ultrasound scanning on the diagnosis and management of fetal abnormalities has also not been demonstrated.

Ultrasound use for fetal scanning is generally considered safe if properly used when information is required about a pregnancy. However, ultrasound is a form of energy and even at low levels, some studies have shown that it can produce physical effects in tissue, such as jarring vibrations and rise in temperature. Although there is no evidence that these physical effects can harm a fetus, the existence of these effects means that prenatal ultrasound cannot be considered completely harmless (U.S. Food and Drug Administration [FDA], 2004). There is increasing concern regarding the use of ultrasound solely for the purpose of providing enhanced photographs and videos of a fetus.

Literature Review: A prospective study (n=118) by Chen et al. (2009) assessed the reproducibility of measurements of nasal bone length using a three-dimensional (3D) ultrasound in the first trimester compared to 2D measurements. The successful rate of measurement of nasal bone length by 3D ultrasound was 79.7%. There was significant inter-method difference between the results obtained by 2D and 3D, substantial variation between observers in 3D measurement of fetal nasal bone length in the first trimester. Independent 3D measurement of nasal bone was found to have no additional advantages over 2D sonography (Chen, et al., 2009).

A prospective randomized controlled study (n=60) by Lapaire et al. (2008) assessed the impact of 3D versus 2D ultrasound on maternal-fetal bonding. Maternal recognition was higher with 3-D US (p=0.004), however the maternal preference of 3D US had no significant impact on maternal-fetal bonding. Another randomized study (n=100) by Rustico et al., (2005) reported that the addition of 4D ultrasound results did not significantly change the perception that women have of their baby nor their antenatal emotional attachment compared with conventional 2D ultrasound.

Randomized controlled and evaluation studies (n=range of 63–3472) comparing the diagnostic accuracy of the different ultrasonographic techniques have found the diagnostic information provided by 3D/4D ultrasound to be consistent with that provided by 2D ultrasound and have reported that 3D ultrasound is most helpful as an adjunct to 2D ultrasound imaging (Lindell and Marsal, 2009; Goncalves, et al., 2006; Merz and Welter, 2005; Michailidis, et al., 2002; Scharf, et al., 2001; Dyson, et al., 2000). Further well designed studies are needed to clearly define the indications for 3D/4D imaging.

Professional Societies/Organizations

According to the ACOG guidelines on ultrasound in pregnancy, the technical advantages of 3D ultrasonography include its ability to acquire and manipulate an infinite number of planes and to display ultrasound planes traditionally inaccessible by 2D ultrasonography. Despite these technical advantages, proof of a clinical advantage of 3D ultrasonography in prenatal diagnosis in general is still lacking. Until clinical evidence shows a clear advantage to conventional 2D ultrasonography, 3D ultrasonography is not considered a required modality at this time (ACOG, 2009).

The American Institute of Ultrasound in Medicine (AIUM) states that 2D sonography is currently the primary method of medically-indicated anatomic imaging with ultrasound. While 3D sonography may be helpful in diagnosis, it should be considered only as a developing technology. Its role is restricted to use as an adjunct only, not as a replacement, for 2D ultrasound. The AIUM also states that the use of either 2D or 3D ultrasound solely to view the fetus, obtain a picture of the fetus, or determine the fetal gender without a medical indication, is inappropriate and contrary to responsible medical practice. Although there are no confirmed biological effects on patients caused by exposures from present diagnostic ultrasound equipment, the possibility exists that such biological effects may be identified in the future (AIUM, 2003, 2005).

The Society of Maternal and Fetal Medicine (SMFM) states that only one medically indicated ultrasound per pregnancy, per practice is appropriate. Once this detailed fetal anatomical exam is done, a second one should not be performed unless there are extenuating circumstances with a new diagnosis (SMFM, 2004).

The National Collaborating Centre for Women's and Children's Health recommends that pregnant women be offered an ultrasound scan to screen for structural anomalies, ideally between 18 and 20 weeks' gestation, by an appropriately trained sonographer and with equipment of an appropriate standard. This scan will also help to

ensure consistency of gestational age assessments, improve the performance of mid-trimester serum screening for Down's syndrome, and reduce the need for induction of labor after 41 weeks. The guideline states that the evidence does not support the routine use of ultrasound scanning after 24 weeks' gestation and therefore it should not be offered (National Collaborating Centre for Women's and Children's Health, 2003).

Summary

Although some controversy still exists regarding whether routine ultrasound screening of all obstetric patients improves pregnancy outcomes, one ultrasound examination per pregnancy is considered the standard of care. Evidence in the published peer-reviewed medical literature as well as professional society opinions support the efficacy of first-trimester ultrasound screening for decreasing post-term labor induction rates. Two-dimensional (2D) ultrasound remains the primary method of obstetric imaging. At present, there is insufficient evidence in the peer-reviewed medical literature to support the use of three-dimensional (3D) or four-dimensional (4D) ultrasound. Although the use of 3D ultrasound is increasing in many clinical settings, the role of this technology is unclear. It has not been demonstrated that any additional information provided by 3D sonography results in improved health outcomes or impacts treatment decisions. There is also insufficient evidence to indicate that the use of 4D ultrasound results in improved fetal outcomes or enhances fetal-parental bonding.

Coding/Billing Information

Note: This list of codes may not be all-inclusive.

Covered when medically necessary:

CPT [®] * Codes	Description
76801	Ultrasound, pregnant uterus, real time with image documentation, fetal and maternal evaluation, first trimester (<14 weeks 0 days), transabdominal approach; single or first gestation
76802	Ultrasound, pregnant uterus, real time with image documentation, fetal and maternal evaluation, first trimester (<14 weeks 0 days), transabdominal approach; each additional gestation (List separately in addition to code for primary procedure)
76805	Ultrasound, pregnant uterus, real time with image documentation, fetal and maternal evaluation, after first trimester (> or = 14 weeks 0 days), transabdominal approach; single or first gestation
76810	Ultrasound, pregnant uterus, real time with image documentation, fetal and maternal evaluation, after first trimester (> or = 14 weeks 0 days), transabdominal approach; each additional gestation (List separately in addition to code for primary procedure)
76811	Ultrasound, pregnant uterus, real time with image documentation, fetal and maternal evaluation plus detailed fetal anatomic examination, transabdominal approach; single or first gestation
76812	Ultrasound, pregnant uterus, real time with image documentation, fetal and maternal evaluation plus detailed fetal anatomic examination, transabdominal approach; each additional gestation (List separately in addition to code for primary procedure)
76815	Ultrasound, pregnant uterus, real time with image documentation, limited (e.g., fetal heart beat, placental location, fetal position and/or qualitative amniotic fluid volume), one or more fetuses
76816	Ultrasound, pregnant uterus, real time with image documentation, follow-up (e.g., re-evaluation of fetal size by measuring standard growth parameters and amniotic fluid volume, re-evaluation of organ system(s) suspected or confirmed to be abnormal on a previous scan), transabdominal approach, per fetus
76817	Ultrasound, pregnant uterus, real time with image documentation, transvaginal
76820	Doppler velocimetry, fetal; umbilical artery
76821	Doppler velocimetry, fetal; middle cerebral artery

ICD-9-CM Diagnosis Codes	Description
622.5	Incompetence of cervix
630	Hydatiform mole
631	Other abnormal product of conception
632	Missed abortion
633.00 - 633.91	Abdominal pregnancy
640.0	Threatened abortion
640.8	Other specified hemorrhage in early pregnancy
640.9	Unspecified hemorrhage in early pregnancy
641.00 – 641.93	Antepartum hemorrhage, abruption placentae, and placenta previa
643.00 – 643.91	Excessive vomiting in pregnancy
644.00 – 644.21	Early or threatened labor
645.1	Post term pregnancy
645.2	Prolonged pregnancy
646.00 – 646.93	Other complications of pregnancy, not elsewhere classified
647.00 – 647.94	Infectious and parasitic conditions in the mother classifiable elsewhere, but complicating pregnancy, childbirth, or the puerperium
648.00 – 648.94	Other current conditions in the mother classifiable elsewhere, but complicating pregnancy, childbirth, or the puerperium
651.00 – 651.93	Multiple gestation
652.00 – 652.93	Malposition and malpresentation of the fetus
654.00 – 654.94	Abnormality of organs and soft tissues of pelvis
655.00 – 655.93	Known or suspected fetal abnormality affecting management of mother
656.00 – 656.93	Other fetal and placental problems affecting management of mother
657.00 – 657.03	Polyhydramnios
658.00 – 658.93	Other problems associated with amniotic cavity and membranes
764.00 - 764.99	Slow fetal growth and fetal malnutrition
766.0 - 766.22	Disorders relating to long gestation and high birthweight
768.0	Fetal death from asphyxia or anoxia before onset of labor or at unspecified time
V22.0 - V22.2	Normal pregnancy
V23.0 - V23.9	Supervision of high risk pregnancy
V28.0 - V28.9	Antenatal screening for abnormalities

Experimental/Investigational/Unproven/Not Covered:

CPT* Codes	Description
76376 [†]	3D rendering with interpretation and reporting of computed tomography, magnetic resonance imaging, ultrasound or other tomographic modality; not requiring image postprocessing on an independent workstation.

76377 [†]	3D rendering with interpretation and reporting of computed tomography, magnetic resonance imaging, ultrasound or other tomographic modality; requiring image postprocessing on an independent workstation.
76499 [†]	Unlisted diagnostic radiographic procedure

†Note: Experimental/Investigational/Unproven and Not Covered when used to report 3D or 4D ultrasound in pregnancy.

ICD-9-CM Diagnosis Codes	Description
	No specific codes

***Current Procedural Terminology (CPT®) © 2010 American Medical Association: Chicago, IL.**

References

1. ACOG Committee on Practice Bulletins-Obstetrics. ACOG Practice Bulletin. Clinical management guidelines for obstetricians-gynecologists. Number 55, September 2004a (replaces practice pattern number 6, October 1997). Management of Postterm Pregnancy. *Obstet Gynecol.* 2004 Sep;104(3):639-46.
2. American College of Obstetricians and Gynecologists. ACOG Practice Bulletin No. 101: Ultrasonography in pregnancy. *Obstet Gynecol.* 2009 Feb;113(2 Pt 1):451-61.
3. American College of Obstetricians and Gynecologists (ACOG) Committee on Ethics. ACOG Committee Opinion. Number 297, August 2004b. Nonmedical use of obstetric ultrasonography. *Obstet Gynecol.* 2004 Aug;104(2):423-4.
4. American Institute of Ultrasound in Medicine, AIUM Practice Guidelines for the Performance of Obstetric Ultrasound Examinations. 2007 Oct. Accessed June 30, 2008. Available at URL address: <http://www.aium.org/publications/clinical/obstetric.pdf>
5. American Institute of Ultrasound in Medicine, Official Statement on 3D technology, 1999. Revised November 12, 2005. Accessed June 30, 2004, July 10, 2006. Available at URL address: <http://www.aium.org/provider/statements/statements.asp>
6. Barnett SB, Maulik D; International Perinatal Doppler Society. Guidelines and recommendations for safe use of Doppler ultrasound in perinatal applications. *J Matern Fetal Med.* 2001 Apr;10(2):75-84.
7. Benacerraf BR, Benson CB, Abuhamad AZ, Copel JA, Abramowicz JS, Devore GR, et al. Three- and 4-dimensional ultrasound in obstetrics and gynecology: proceedings of the american institute of ultrasound in medicine consensus conference. *J Ultrasound Med.* 2005 Dec;24(12):1587-97.
8. Bennett KA, Crane JM, O'Shea P, Lacelle J, Hutchens D, Copel JA. First trimester ultrasound screening is effective in reducing post-term labor induction rates: A randomized controlled trial. *Am J Obstet Gynecol.* 2004 Apr;190(4):1077-81.
9. Bricker L, Garcia J, Henderson J, et al. Ultrasound screening in pregnancy: A systematic review of the clinical effectiveness, cost-effectiveness and women's views. *Health Technol Assess.* 2000;4(16):i-vi, 1-193.
10. Chen M, Wang HF, Leung TY, Fung TY, Chan LW, Sahota DS, et al. First trimester measurements of nasal bone length using three-dimensional ultrasound. *Prenat Diagn.* 2009 Aug;29(8):766-70.

11. Dyson RL, Pretorius DH, Budorick NE, Johnson DD, Sklansky MS, Cantrell CJ, et al. Three-dimensional ultrasound in the evaluation of fetal anomalies. *Ultrasound Obstet Gynecol.* 2000 Sep;16(4):321-8.
12. ECRI Institute. *Ultrasound Screening of Normal-Risk Women for Prevention of Postterm Pregnancy.* Plymouth Meeting (PA): ECRI Institute Health Technology Assessment Information Service; 2003 September. 67 p. (Evidence Report; no. 147). Available at URL address: <http://www.ecri.org>
13. Goncalves LF, Lee W, Espinoza J, Romero R. Three- and 4-dimensional ultrasound in obstetric practice: does it help? *J Ultrasound Med.* 2005 Dec;24(12):1599-624.
14. Goncalves LF, Nien JK, Espinoza J, Kusanovic JP, Lee W, Swope B, et al. What does 2-dimensional imaging add to 3- and 4-dimensional obstetric ultrasonography? *J Ultrasound Med.* 2006 Jun;25(6):691-9.
15. Institute for Clinical Systems Improvement (ICSI). *Health Care Guideline: Routine Prenatal Care.* August 2009. Accessed July 11, 2010. Available at URL address: http://www.icsi.org/prenatal_care_4/prenatal_care__routine__full_version__2.html
16. Lapaire O, Alder J, Peukert R, Holzgreve W, Tercanli S. Two- versus three-dimensional ultrasound in the second and third trimester of pregnancy: impact on recognition and maternal-fetal bonding. A prospective pilot study. *Arch Gynecol Obstet.* 2007 Nov;276(5):475-9. Epub 2007 Apr 25.
17. Merz E, Welter C. 2D and 3D Ultrasound in the evaluation of normal and abnormal fetal anatomy in the second and third trimesters in a level III center. *Ultraschall Med.* 2005 Feb;26(1):9-16.
18. Michailidis GD, Papageorgiou P, Economides DL. Assessment of fetal anatomy in the first trimester using two- and three-dimensional ultrasound. *BR J Radiol.* 2002 Mar;75(891):215-9.
19. National Collaborating Centre for Women's and Children's Health. *Antenatal care: routine care for the healthy pregnant woman.* London: RCOG Press; 2003 Oct. 286 p. Accessed July 10, 2006. Available at URL address: http://www.guideline.gov/summary/summary.aspx?doc_id=4808&nbr=003470&string=antenatal+AND+care
20. National Institutes of Health (NIH). *NIH Consensus Statement. Diagnostic Ultrasound Imaging in Pregnancy.* Accessed July 8, 2004. Available at URL address: http://www.consensus.nih.gov/cons/041/041_statement.htm
21. Rustico MA, Mastromatteo C, Grigio M, Maggioni C, Gregori D, Nicolini U. Two-dimensional vs. two-plus four-dimensional ultrasound in pregnancy and the effect on maternal emotional status: a randomized study. *Ultrasound Obstet Gynecol.* 2005 May;25(5):468-72.
22. Scharf A, Ghazwiny MF, Steinborn A, Baier P, Sohn C. Evaluation of two-dimensional versus three-dimensional ultrasound in obstetric diagnostics: a prospective study. *Fetal Diagn Ther.* 2001 Nov-Dec;16(6):333-41.
23. Society for Maternal-Fetal Medicine (SMFM), Coding Committee. *White paper on ultrasound code 76811. Announcements.* Washington, DC: SMFM; May 24, 2004. Accessed July 8, 2005. Available at URL address: <http://www.smfm.org/index.cfm?zone=news&nav=viewnews&newsID=238&smfm=on=yes>.
24. Timor-Tritsch IE, Monteagudo A. Three and four-dimensional ultrasound in obstetrics and gynecology. *Curr Opin Obstet Gynecol.* 2007 Apr;19(2):157-75.
25. U.S. Food and Drug Administration, FDA Consumer magazine. *FDA cautions against ultrasound 'keepsake' images.* January- February 2004. Accessed July 8, 2005. Available at URL address: http://www.fda.gov/fdac/features/2004/104_images.html

26. Whitworth M, Bricker L, Neilson JP, Dowswell T. Ultrasound for fetal assessment in early pregnancy. Cochrane Database Syst Rev. 2010 Apr 14;4:CD007058.

Policy History

<u>Pre-Merger Organizations</u>	<u>Last Review Date</u>	<u>Policy Number</u>	<u>Title</u>
CIGNA HealthCare	8/15/2008	0142	Ultrasound In Pregnancy (including 3D and 4D Ultrasound)

"CIGNA" and the "Tree of Life" logo are registered service marks of CIGNA Intellectual Property, Inc., licensed for use by CIGNA Corporation and its operating subsidiaries. All products and services are provided exclusively by such operating subsidiaries and not by CIGNA Corporation. Such operating subsidiaries include Connecticut General Life Insurance Company, CIGNA Behavioral Health, Inc., Intracorp, and HMO or service company subsidiaries of CIGNA Health Corporation and CIGNA Dental Health, Inc. In Arizona, HMO plans are offered by CIGNA HealthCare of Arizona, Inc. In California, HMO plans are offered by CIGNA HealthCare of California, Inc. and Great-West Healthcare of California, Inc. In Connecticut, HMO plans are offered by CIGNA HealthCare of Connecticut, Inc. In North Carolina, HMO plans are offered by CIGNA HealthCare of North Carolina, Inc. In Virginia, HMO plans are offered by CIGNA HealthCare Mid-Atlantic, Inc. All other medical plans in these states are insured or administered by Connecticut General Life Insurance Company.

Connecticut General Life Insurance Company has acquired the business of Great-West Healthcare from Great-West Life & Annuity Insurance Company (GWLA). Certain products continue to be provided by GWLA (Life, Accident and Disability, and Excess Loss). GWLA is not licensed to do business in New York. In New York, these products are sold by GWLA's subsidiary, First Great-West Life & Annuity Insurance Company, White Plains, N.Y.