



CIGNA MEDICAL COVERAGE POLICY

The following Coverage Policy applies to all health benefit plans administered by CIGNA Companies including plans formerly administered by Great-West Healthcare, which is now a part of CIGNA.

Effective Date 5/15/2011
Next Review Date 11/15/2011
Coverage Policy Number 0234

Subject **Varicose Vein Treatments**

Table of Contents

Coverage Policy	1
General Background	4
Coding/Billing Information	12
References	16
Policy History	24

Hyperlink to Related Coverage Policies

INSTRUCTIONS FOR USE

Coverage Policies are intended to provide guidance in interpreting certain **standard** CIGNA HealthCare benefit plans. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement (GSA), Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document **always supercedes** the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. Proprietary information of CIGNA. Copyright ©2011 CIGNA

Coverage Policy

Coverage for treatment of varicose veins is dependent on benefit plan language and may be subject to the provisions of a cosmetic and/or reconstructive surgery benefit. Under many benefit plans, treatment of varicose veins is not covered when provided solely for the purpose of altering appearance or self-esteem or to treat psychological symptomatology or psychosocial complaints related to one's appearance. In addition, some benefit plans specifically exclude coverage for the invasive treatment of varicose veins. Please refer to the applicable benefit plan document to determine benefit availability and the terms, conditions and limitations of coverage.

If coverage is available for the treatment of varicose veins, the following conditions of coverage apply.

Invasive varicose vein treatments that may be considered for coverage include the following modalities, as a single or combined treatment, when the specific criteria for the procedure(s) outlined below are met:

- ambulatory phlebectomy
- ligation and excision
- radiofrequency ablation (RFA)
- endovenous laser therapy (EVLT)
- sclerotherapy (liquid, foam, ultrasound-guided, or endovenous chemical ablation)

COVERAGE CRITERIA FOR HIGH RISK INDICATIONS

CIGNA covers ambulatory phlebectomy, ligation and excision, RFA, EVLT, and/or sclerotherapy (i.e., liquid, foam, ultrasound-guided, endovenous chemical ablation) as medically necessary for ANY of the following HIGH RISK varicose vein indications:

- leg ulceration(s) due to saphenous vein insufficiency refractory to conservative management
- recurrent bleeding from the saphenous vein or other varicosity
- history of a significant episode of bleeding from a varicosity

COVERAGE CRITERIA FOR LOWER RISK INDICATIONS

CIGNA covers ambulatory phlebectomy, ligation and excision, RFA, and/or EVLT for the treatment of symptomatic saphenous varicose veins as medically necessary for ANY of the following indications:

- pain resulting in a clinically significant functional impairment (e.g., inability to perform household chores or prolonged standing, interference with essential job functions)
- recurrent phlebitis or thrombophlebitis
- refractory dependent edema
- persistent stasis dermatitis
- chronic cellulitis

when ALL of the following criteria are met:

- a Doppler and/or Duplex ultrasonography evaluation and report, performed no more than 12 months prior to the requested procedure, confirms incompetence/reflux and documents vein size ≥ 3 mm
- documentation of **BOTH** of the following:
 - previous invasive treatment(s) of varicose veins (if any)
 - failure or intolerance of medically supervised conservative management, including but not limited to compression stocking therapy for three consecutive months
- a clearly defined treatment plan including the procedure (CPT) codes for the planned interventions and whether the proposed treatment is to the left leg, the right leg, or both legs

CIGNA covers sclerotherapy* (liquid, foam, ultrasound-guided, or endovenous chemical ablation) as medically necessary treatment of symptomatic saphenous varicose veins or tributaries when BOTH of the following criteria are met:

- reflux at the saphenofemoral or saphenopopliteal junction (i.e., proximal to the incompetence) is concurrently being/has previously been treated
- **EITHER** of the following:
 - Sclerotherapy* is performed as an adjunctive treatment to one of the above listed modalities on the same date of service
 - Sclerotherapy* is performed for treatment of residual or recurrent varicosities, when performed within 12 months of a prior invasive varicose vein procedure

***Note: Coverage for sclerotherapy for these indications is limited to a maximum of three (3) sclerotherapy treatment sessions per leg, without additional clinical documentation, when performed within 12 months of the initial invasive varicose vein procedure.**

COVERAGE CRITERIA FOR ADDITIONAL SCLEROTHERAPY TREATMENT SESSION(S)

Requests for additional sclerotherapy treatment, extending beyond the maximum three treatment sessions per leg, may be considered for coverage when ALL of the following additional criteria have been met:

- additional documentation confirms persistence of symptoms despite prior invasive treatment
- Doppler or Duplex reports and/or standing photographs (when appropriate) confirm persistent veins greater than 3 mm in size
- evidence of a clearly defined treatment plan including the procedure codes for the planned intervention

COVERAGE CRITERIA FOR SEPS

CIGNA covers subfascial endoscopic perforator surgery (SEPS) as medically necessary when ALL of the following medical necessity criteria are met:

- a Doppler and/or Duplex ultrasonography evaluation and report, performed no more than 12 months prior to the requested procedure, confirms reflux of the incompetent perforator vein and location on the medial aspect of the calf being treated
- failure or intolerance of medically supervised conservative management, including but not limited to compression stocking therapy, for at least three consecutive months
- documentation of at least **ONE** of the following conditions:
 - venous stasis dermatitis/ulceration
 - chronic venous insufficiency

COVERAGE CRITERIA FOR ADDITIONAL SCLEROTHERAPY TREATMENT SESSION

Requests for additional sclerotherapy treatment, extending beyond the maximum three treatment sessions per leg, may be considered for coverage when ALL of the following additional criteria have been met:

- additional documentation confirms persistence of symptoms despite prior invasive treatment
- Doppler or Duplex reports and/or standing photographs (when appropriate) confirm persistent veins greater than 3 mm in size
- evidence of a clearly defined treatment plan including the procedure codes for the planned intervention

Requests for treatment sessions extending beyond one year from the initial invasive treatment session will be similarly subject to a new medical necessity review, including submission of all the materials normally required for an initial submission.

NOT COVERED

CIGNA does not cover ANY of the following varicose vein treatments because each is considered cosmetic in nature as well as not medically necessary:

- treatment of telangiectasis or varicose veins that are less than 3 mm in diameter by any method
- sclerotherapy with glycerin/glycerol
- intense pulsed-light source (photothermal sclerosis) treatment of a varicose vein

CIGNA does not cover ANY of the following varicose vein treatments, because each is considered experimental, investigational or unproven (this list may not be all-inclusive):

- non-compressive sclerotherapy
- transdermal laser therapy
- transilluminated powered phlebectomy (TIPP, TriVex™)
- SEPS for the treatment of venous insufficiency as a result of post-thrombotic syndrome
- sclerotherapy (i.e., liquid, foam, ultra-sound guided, endovenous chemical ablation) when performed for ANY of the following indications:
 - sole treatment of accessory, reticular or varicose tributaries without associated occlusion of the saphenofemoral or saphenopopliteal junction
 - incompetence that is isolated to the perforator veins
 - of the Greater Saphenous Vein (GSV), with or without associated ligation of the saphenofemoral junction
 - as a sole (i.e., stand alone) treatment for reflux occurring at the saphenofemoral or saphenopopliteal junction
- endomechanical ablative approach (e.g., ClariVein™ Catheter)
- cryostripping, (including cryoablation, cryofreezing) of any vein

General Background

Varicose veins result from weakening or incompetence of a one-way valve, leading to a retrograde flow or reflux of blood in the vessel. The varicosity may vary in size from 3–10 mm on average. Symptoms that have been reported as associated with varicose veins of the lower extremities include pain, cramping, aching, burning, throbbing, swelling and the feeling of heaviness or fatigue in the leg. Typically, symptoms are exacerbated by standing and warm weather (Hamper, et al., 2007). Saphenous varicose veins can ultimately result in intractable ulcerations and recurrent bleeding. Patients with larger varicosities (e.g., varicose veins greater than 3 mm in diameter) are more prone to thrombophlebitis and other complications than those with smaller varicosities. Chronic cellulitis may also be associated with varicosities.

The venous system of the lower extremities is separated into two main systems: the deep venous and the superficial venous system. The two systems are connected by perforator veins. The deep venous system comprises the popliteal and femoral veins; the superficial venous system comprises the greater saphenous and short saphenous veins (formerly called the lesser saphenous vein). The GSV generally measures 3–4 mm in diameter in the upper thigh; the GSV meets the femoral vein at the saphenofemoral junction (SFJ). Approximately 60% of patients who have varicosities have reflux in the GSV (Hamper, et al., 2007). The short saphenous vein is not usually larger than 3 mm in diameter, and connects with the deep veins at the saphenopopliteal junction (SPJ) in the knee area. Incompetence of the superficial venous system typically results from failure of valves at the SFJ and the SPJ with resulting pressure that is worse at the more distal area of the vein. Incompetence of the perforating veins also leads to increased pressure in the superficial venous system due to the pump mechanism of the calf. Varicose tributaries are veins that empty into a larger vein.

Telangiectases are permanently dilated blood vessels, also called spider veins that create fine red or blue lines on the skin. They are similar to varicose veins, but are limited to the dermis and are not usually more than 3 mm in diameter. They are not typically associated with symptoms, and treatment is generally considered cosmetic in nature and not medically necessary.

Varicose veins may develop during pregnancy, although surgery or sclerotherapy is not typically performed, as the treatment is not medically necessary. Most varicosities will spontaneously resolve within 4–6 months after delivery.

Varicose veins of the upper extremity are rare; still there are a few reports in the published, peer-reviewed medical literature dealing with the management of upper extremity varicosities (Welch and Villavicencio, 1994; Duffy, et al., 1999; Lee, 2002; Bowes and Goldman, 2002). However, authors have reported successful outcomes utilizing methods of treatment similar to lower extremity varicosities (e.g., sclerotherapy, ligation and stripping, phlebectomy).

Various ultrasound technologies are used in conjunction with other noninvasive testing to determine the physiological characteristics of the varicosities, as physical exam alone may not be reliable. Duplex ultrasound, Doppler ultrasound and plethysmography may all be used to diagnose varicose veins. In most cases, once the initial vein mapping is performed, it is not essential that follow-up scanning be done for subsequent sclerotherapy sessions. It has not been demonstrated in the published medical literature that repeat Duplex or Doppler studies are essential for the successful outcome of the procedure when performed as part of a series of sclerotherapy sessions. Also, routine use of any of these tools in the absence of venous symptoms or clinical evidence of venous insufficiency or reflux is not considered a medical necessity. Photographs or diagrams are helpful in assessing the size and extent of the varicosities.

The CEAP classification is a method commonly used to document the severity of chronic venous disease and is based on clinical presentation (C), etiology (E), anatomy (A), and pathophysiology (P) (See Table 1).

Table 1: CEAP Classification

Class	Definition
C - Clinical Classification , supplemented by “A” for asymptomatic and “S” for symptomatic presentation	Class 0: No visible or palpable signs of venous disease Class 1: Telangiectasia, reticular veins, malleolar flare Class 2: Varicose veins Class 3: Edema without skin changes Class 4: Skin changes ascribed to venous disease (e.g., pigmentation, venous eczema, lipodermatosclerosis) Class 5: Skin changes as defined above with healed ulceration Class 6: Skin changes as defined above with active ulceration
E - Etiology	Congenital, Primary, Secondary, No venous disease
A - Anatomy	Superficial, Perforator, Deep, No venous location
P - Pathophysiology	Reflux or obstruction (alone or combined); Basic or Advanced

Classification of disease starts with an initial assessment and may not be entirely completed until after surgery and histopathologic assessment. As a result, it is recommended that CEAP classification value be followed by the date of examination. Venous disease can be reclassified at any given time. It is also recommended that the level of investigation be included, with Level I representing the office visit, Level II representing noninvasive venous laboratory testing and Level III representing invasive assessment and more complex imaging studies.

Various methods of treatment have been investigated. Conservative medical practices that may be used in the management of varicose veins include leg elevation, analgesia for symptom relief and avoidance of prolonged periods of standing. Compression therapy, the use of custom-fit compression stockings with pressure gradients, a mainstay of initial/conservative management, is routinely attempted prior to stripping, ligation, sclerotherapy or other, more invasive procedures. The amount of compression required for treatment of stasis dermatitis or ulceration is between 35 and 40 mm Hg, for varicose veins, for mild edema and leg fatigue the recommended pressure is 20 to 30 mm Hg (Habif, 2009). When conservative measures fail, treatment options rely on identifying and correcting the site of reflux and on redirecting the flow of blood through veins with properly functioning valves. No single method of treatment is universally employed in the literature; the intervention selected is generally dependent upon the competency of deep and perforating veins, and the site and degree of reflux. Surgery is commonly used to treat mainstem varicose veins. Many patients require a combination of techniques to correct symptoms associated with venous insufficiency, most of which can be performed in a single treatment session. Endovenous thermal ablation procedures include radiofrequency ablation (RFA) and endovenous laser therapy (EVLT). While staging of procedures is generally not required, repeat sclerotherapy sessions may be required for an unsuccessful vein occlusion. Typically, a treatment plan includes thermal ablation of the incompetent vein segment for greater saphenous vein reflux and associated larger varicosities. Once this segment is treated, if there are associated varicosities greater than 4mm in diameter phlebectomy is often performed during the same session, and for smaller veins, sclerotherapy is employed as the treatment of choice (Kouri, 2009).

Complications associated with varicose vein treatment vary and are dependent on the type of treatment employed. Complications that may result from sclerotherapy and phlebectomy include but are not limited to hyperpigmentation, allergic skin reactions, migraine-like symptoms (particularly from foam sclerosants), pain at the injection site, superficial and deep thromboembolic events and subcutaneous hematomas. Most complications are transient and resolve with conservative measures. Subcutaneous hematoma formation is easily managed with warm compresses and nonsteroidal anti-inflammatory medications. Thromboembolic events although rare can be life-threatening and may require anticoagulation (Lew, Weaver, 2010; Alaiti, 2010). Complications associated with thermal ablation techniques are usually minor and self limiting; serious events are rare.

Invasive Approaches

Sclerotherapy: Sclerotherapy is an invasive procedure used to eradicate small to medium sized varicose veins of the superficial venous system (greater and small saphenous veins). When reflux is present at the junction, sclerotherapy should be performed in addition to surgical ligation and division of the junction, promoting control of the point of reflux. Injection of the vein at its junction and of the incompetent perforating veins has been

proposed as an alternative to ligation, however, the scientific literature does not support the efficacy of this procedure. Sclerotherapy has not been shown to be effective as a sole treatment of larger incompetent veins and is often used with other approaches to treat significant varicosities. According to the American Academy of Cosmetic Surgery (AACS) guidelines for sclerotherapy (2003) effectiveness approaches 90-95% when sclerotherapy is used for the treatment of small diameter vessels, for vessels larger than 4mm in diameter treatment failures average 25%, with recurrences noted in 10% of large diameter vessels. A recent systematic review (Leopardi, et al., 2010) also supports sclerotherapy is indicated for patients with minor, superficial veins not related to reflux in the saphenous system. Corabian et al. (2004) noted the role of sclerotherapy in the management of GSV and perforator incompetence has not been clearly defined, although sclerotherapy may be indicated for treatment of large saphenous veins without reflux (Corabian, et al., 2004).

During sclerotherapy, the abnormal vein is injected with a sclerosing agent that irritates the lining of the vein, causing it to thrombose and stenose, ultimately leading to resorption into the surrounding tissue. Echosclerotherapy using either liquid or foam sclerosant, also referred to as ultrasound-guided sclerotherapy and endovenous chemical ablation (ECA), employs real-time ultrasound during the sclerotherapy procedure to help locate deep or inaccessible sites. Echosclerotherapy is indicated for treatment of veins below the surface, such as deep veins and other varices that are difficult to visualize (Corabian, et al., 2004). According to the ACP, (2008) the use of image guided techniques such as ultrasound is essential for the safe and effective performance of endovenous chemical ablation and reflects the current standard of care.

Foam sclerotherapy, which involves the use of a sclerosing solution that has been forcibly mixed with air or gas (e.g., carbon dioxide) to create a foam agent, is often used in large-diameter vessels and with the use of ultrasound. Ultrasound is used to monitor the foam distribution. Foam sclerosant forces blood out of the vein and allows for less dilution of the sclerosant and more contact with the endothelium (Lew, Weaver, 2010). Overall, authors generally agree foam sclerotherapy is a safe and effective method of treating varicose veins (Rabe, et al., 2004; Wright, et al., 2006; Kendler, et al., 2007; Uurto, et al., 2007; Subramonia and Lees, 2007; Jia, et al., 2007; Darvall, et al., 2009). In addition, this method is supported by several professional societies and organizations as being safe and at least equally if not more effective than liquid sclerotherapy (American Academy of Cosmetic Surgery [AACS] 2003; National Institute for Clinical Excellence [NICE], 2007; American College of Phlebology, 2008; German Society of Phlebology [Rabe, Pannier, 2010]).

As with sclerotherapy in general, the need for repeat treatment sessions when utilizing any of these methods of treatment has been reported in the literature (Barrett, et al., 2004; Darke, and Baker, 2006). Although echosclerotherapy has been investigated as an alternative to traditional saphenous vein ligation and stripping (Min, Navarro, 2000; Bountouroglou, et al., 2006), there is insufficient evidence in the medical literature to support safety, efficacy and improvement in long-term clinical outcomes when used for this indication. Evidence consists mainly of case series with few comparative trials and mixed reported clinical outcomes.

There is no consensus in the published scientific literature regarding the optimal number of sclerotherapy treatments required to reduce the symptoms associated with varicose veins and the number treatments needed to resolve symptoms varies among patients. The AACS (2003) reports sclerotherapy is the treatment of choice for varicose veins that are 2–4 mm in diameter and large areas of veins can usually be eradicated using two to three treatment sessions. Vessels 4–6 mm in diameter may be treated by sclerotherapy or ambulatory phlebectomy. Weiss et al. (1992) reported that, in some cases, four or more separate sclerosing treatments may be necessary to completely eradicate groups of varicose veins; such a course of treatment might include 1–4 treatments for a region of the leg or three treatments for a larger vein coursing several regions of the leg.

The primary aims of sclerotherapy are to prevent complications of varicose disease and relieve symptoms; cosmetic improvement in the leg's appearance is an added benefit. Treatment provided solely for cosmetic purposes is not considered a medical necessity. Sclerotherapy is a palliative solution and cannot prevent the formation of new varicosities. New varicosities may form, either because of an underlying illness or condition, or, in some cases, because of a genetic predisposition.

In compressive sclerotherapy, the most commonly performed method of sclerotherapy, compressive dressings are applied after injection of the sclerosing agent, while the limb is elevated and the vein is drained. External compression and internal decompression (e.g., walking) stimulates fibrosis, which contributes to obliteration of the entire vein wall (Labas, et al., 2003). Non-compressive sclerotherapy involves injecting a sclerosant into the

non-elevated (blood-filled) vein without applying a compressive dressing. This method of therapy has not been shown to be effective in producing long-term obliteration of the incompetent veins.

Sclerosing agents currently approved by the U.S. Food and Drug Administration (FDA) to treat varicose veins of the lower extremities and most commonly used include sodium tetradecyl sulfate (Sotradecol[®]) and polidocanol (Asclera[®]); polidocanol was approved by the FDA March 2010 for the treatment of small spider veins and reticular veins. According to the manufacturer Asclera has not been studied in varicose veins larger than 3mm. Other agents such as morrhuate sodium (Scleromate[™] morrhuate sodium) although FDA approved are not used as commonly. Glycerin/ glycerol is an osmotic dehydrating agent which is primarily used for the treatment of residual telangiectasias (Duffy, 2010). Nonetheless, there is no evidence-based consensus on the optimal type, dosage or concentration of the sclerosing agent.

Transdermal Light/Laser Therapy: Photothermal sclerosis, such as PhotoDerm[®] Vasculite[™], is also referred to as intense pulsed-light source. Used as an alternative to or to complement sclerotherapy in treating small varicose veins and telangiectasias (spider veins), this type of light therapy utilizes small pulses of light energy which travel through the skin, are absorbed by the blood, are then changed to heat and ultimately destroy the vein. Successful treatment requires adequate heating of the veins, and several treatments are usually required for optimal results.

Transcutaneous laser ablation, also known as transdermal laser treatment, is a type of laser therapy similar to light therapy that involves the use of a laser to treat small varicose and spider veins. Small laser pulses are delivered to the vein, causing heat, which will ultimately lead to destruction of the vein. This modality is not generally useful as a primary treatment of spider veins of the lower extremity; instead, it is employed to treat superficial vessels on the face. The treatment may result in superficial skin burns and permanent pigmentation changes.

Laser or light therapy has been indicated for the treatment of telangiectasis and cutaneous vascular lesions (Raulin, et al., 1997; Angermeier, 1999). However, evidence in the published scientific literature indicates that transdermal light/laser therapy has not been shown to be as effective for the lower extremities as for facial telangiectasis and smaller varicosities (Weiss, Dover, 2002). The vessels in the lower extremities are located deeper and have thicker surrounding tissue. Deeper vessels require a longer wavelength and longer pulse duration to damage the vessel effectively. Additionally, because spider veins and varicosities smaller than 3 mm do not usually cause symptoms, they are considered cosmetic; hence, treatment for them is not medically necessary.

Ligation, Division and/or Excision: The traditional surgical treatment of saphenous-vein varicosities consists of surgical ligation and stripping. When the GSV and SSV have reflux or incompetence, junction ligation with or without vein stripping is often recommended; in most cases, ligation is followed by GSV stripping. During the procedure, the saphenous vein and other smaller veins are exposed through an incision in the groin, where the veins are then ligated (i.e., tied off) with sutures. A second incision is made just below the knee or at the ankle to allow access for stripping the vein. When both ends of the vein are free, a wire-like instrument is threaded through the vein, extending up to the second incision in the groin area. The vein is then pulled (i.e., stripped) and removed from the leg. Removal of the superficial symptomatic vein restores venous circulation and provides relief of symptoms. Operative excision of the vein is most often reserved for large varicosities and for those located in the medial or anterior thigh.

Cryostripping: Cryoablation uses extreme cold to cause injury to the vessel. Cryostripping of the GSV may be considered an alternative approach to traditional ligation and stripping. During this procedure, a cryoprobe is passed through the GSV, the probe freeze attaches to the GSV and stripping is performed by pulling back the probe. In one randomized clinical trial (n=494) comparing cryostripping with conventional stripping of the GSV (Klem, et al., 2009) the authors reported that cryostripping accounted for higher failures and residual GSV and offered no benefits over conventional stripping. Menyhei et al. (2008) compared conventional stripping and cryostripping and assessed quality of life outcomes and complications (n=160) in a randomized trial. The authors reported significantly improved quality of life scores for both groups, with no difference between the two groups at six months. There was less bruising in the cryo group but no difference in post-operative pain scores between the two groups. The results of another randomized trial (n=120) indicated that EVLT and cryostripping were similarly effective at two years follow-up (recurrent incompetence 77% and 66%, for EVLT and cryostripping, respectively), however EVLT was superior with regard to duration of operation, postprocedural

pain, induration and resumption of normal activity (Disselhoff, et al., 2008). The published evidence is mixed and does not lend strong support to improved clinical outcomes when compared to more conventional methods of varicose vein treatment. Further studies are needed to demonstrate safety, efficacy and the clinical utility of cryostripping.

Ambulatory Phlebectomy/Stab Phlebectomy: Ambulatory phlebectomy is also widely accepted as an alternative to sclerotherapy, performed alone or in addition to stripping and ligation for the treatment of surface varicose veins. It is also referred to as miniphlebectomy or stab avulsion. In ambulatory phlebectomy, multiple small incisions are made, and the varicose veins are grasped with a small hook or hemostat. They are then clamped, divided and finally extracted. The entire varicosity can be extracted with multiple small incisions. Compression therapy has been shown to reduce bleeding and improve resorption following this method of treatment and is thus widely used for that purpose. Effectiveness is dependent on the type of vein treated; the results of a recent systematic review (Leopardi, et al., (2010) indicated that phlebectomy appears to be a treatment of choice for smaller veins such as the lateral accessory veins, and that for larger veins such as the saphenous veins, phlebectomy may not provide the same level of success as sclerotherapy.

Transilluminated Powered Phlebectomy (TIPP): TIPP, which is similar to ambulatory phlebectomy, is another minimally invasive alternative to standard surgery for the treatment of symptomatic varicosities. Also known as the TriVex™ (Smith & Nephew Inc., Andover, MA) procedure, TIPP involves endoscopic resection and ablation of the superficial varicosity.

Subcutaneous transillumination and tumescent anesthesia help visualize and locate the varicosity, while subcutaneous vein ablation is performed using a powered resector to obliterate the vein. Tumescent anesthesia involves the infusion of large amounts of saline and lidocaine to reduce hemorrhage and of epinephrine to delay absorption of the lidocaine. During this procedure, the veins are marked with a marker, and a bright light is introduced into the leg through a small incision (2–3 cm) to enhance visualization of the veins. The power vein resector is then inserted to cut and remove the vein through suction.

Proponents of this method suggest that the illuminating light allows quicker and more accurate removal of the vein, leading to a more effective yet less traumatic procedure. TIPP is intended for patients who are suitable candidates for conventional ambulatory phlebectomy, and may also be used as an adjunctive method to other varicose vein treatments (e.g., ligation and stripping).

The individual components of the TriVex system were approved for use by the FDA in 1999, however since that time, several other illumination and powered-resection devices have been approved and are available for use.

Evidence evaluating TIPP for the treatment of varicose veins is primarily in the form of published reviews, comparative trials (few involving randomized groups) and both retrospective and prospective case series involving small populations and evaluating short-term outcomes (Franz and Knapp, 2008; Passman, et al., 2007; Scavee, 2006; Chetter, et al., 2006; Aremu, et al., 2004; Shamiyeh, et al., 2003; Scavee, et al., 2003; Chesire, et al., 2002; Spitz, et al., 2000). Two controlled studies specifically compared TIPP to phlebectomy (Aremu, et al., 2004; Scavee, et al., 2003), although neither of these studies were blinded. In addition, the outcomes measured in most studies include operative time, number of incisions, complications, and cosmetic satisfaction with few patient-oriented outcomes being reported. Generally, the results of these studies demonstrate that TIPP is associated with fewer incisions (Luebke, et al., 2008; Chetter, et al., 2006; Aremu, et al., 2004; Shamiyeh, et al., 2003; Scavee, et al., 2003; Spitz, et al., 2000). Operative time varies among authors and with experience. Despite reports in the published literature of a reduced number of incisions, an increase in bruising, postoperative pain and decreased quality of life during the early postoperative period has been reported. Moreover, it has been reported in the literature that technical complications may be associated with inexperience. The published, peer-reviewed, scientific literature does not lead to strong conclusions that TIPP results in clinical outcomes (e.g., improved pain, less varicose vein recurrence) that are as good as treatment with standard conventional methods (i.e., hook phlebectomy). Furthermore, long-term safety and efficacy of the procedure has not been adequately demonstrated.

ECRI Institute published an emerging technology report (2008) evaluating TIPP for treatment of varicose veins. According to the report, the available data are promising for demonstrating the safety and efficacy of TIPP relative to hook phlebectomy and stab avulsion to treat varicose veins. However, ECRI also reported that the available evidence is inadequate to draw firm conclusions about its relative short- and long-term effectiveness,

or its purported advantages over existing methods in terms of complications, operating time, pain, varicose vein recurrence, and cosmetic outcomes.

In 2004 NICE issued an Interventional Procedure Guidance for TIPP. The advisory committee indicated that, although the evidence suggested that the procedure is effective, the data are too limited to be conclusive. In addition, there are no long-term follow-up data (NICE, 2004a).

Endoluminal Radiofrequency Ablation (RFA): Radiofrequency ablation, also known as endovascular occlusion, is a treatment for symptomatic varicose veins that involves delivery of controlled radiofrequency (RF) energy through a catheter inserted into the affected vein. The heat generated by the RF energy causes the vein to contract and become occluded. The treatment is intended as a minimally-invasive alternative to standard surgery for symptomatic varicosities located mainly below the saphenofemoral or saphenopopliteal junction—RFA has also been investigated as a treatment of incompetent perforator veins (Singh and Sura, 2008; Uchino, 2007; Roth, et al. 2007; Peden and Lumsden, 2007; Gibson, et al, 2007a), however data supporting safety and efficacy is limited and further clinical studies are needed to support widespread use for this indication.

RFA using the VNUS® Closure System is a three-part procedure that begins with imaging of the greater saphenous vein, followed by the administration of anesthesia between the vein and the skin. Next, the closure catheter is inserted into the vein, and electrodes are implanted in the venous wall. RF energy is released until the venous wall temperature reaches approximately 85 °C. The temperature is maintained for 30 seconds; then the catheter is slowly retracted, causing the entire length of the vein to collapse on it. If the assessment following treatment indicates any areas of steady flow, those areas may be re-treated, as long as the catheter is reinserted immediately (Chandler, et al., 2000; VNUS, 2000). Possible complications include vessel perforation, pulmonary embolism, phlebitis, hematoma, infection, paresthesia and skin burns (Chandler, et al., 2000; Goldman, 2000; VNUS, 2000).

Evidence in the peer-reviewed published scientific literature supports the safety and efficacy of RFA for the treatment of symptomatic varicose veins. Most early studies were small case series with short-term follow-up (Ogawa, et al., 2005; Goldman, 2002; Weiss, 2002; Goldman, 2000), and only two included direct comparisons with standard treatments (Lurie, 2003; Rautio, 2002). RFA has been shown in a prospective nonrandomized trial to be more effective than foam sclerotherapy for closure of the GSV at one year follow-up (Gonzalez-Zeh, et al., 2008). More recently, RFA has been compared to procedures such as EVLT (Almeida, et al., 2009) and has been evaluated with and without ligation of the saphenofemoral junction (Disselhoff, et al, 2008) in randomized controlled trials. Compared to EVLT, at one month following treatment, RFA was significantly superior for measures evaluating post procedure recovery and quality of life parameters. When performed with and without ligation, at two years post procedure, there was no difference in outcomes (recurrence, degree of ablation and venous clinical severity scores) from adding the ligation procedure. The short-term results of several other studies have demonstrated that the procedure effectively occludes incompetent veins following RFA treatment (Merchant and Pichot, 2006; Hinchliffe et al., 2006; Welch, 2006; Lurie, et al., 2005). Long-term occlusion rates were reported by Merchant and Pichot (2005). This group of authors collected data to evaluate the long-term treatment outcomes of endovascular RFA and to determine risk factors that affect treatment efficacy. In their study, the authors reported on five-year follow-up results of 1006 patients (1222 limbs) treated with radiofrequency obliteration (RFO). Immediate vein occlusion was achieved in 96.8% of limbs confirmed by Duplex ultrasound examination one week or less after the procedure. The vein occlusion rate at six months, one, two, three, four and five years was 89.2%, 87.1%, 88.2%, 83.5%, 84.9% and 87.2%, respectively. The absence of reflux rate was 91.3%, 88.2%, 88.2%, 88.0%, 86.6% and 83.8%, respectively. Over a five-year follow-up period, anatomical failure was identified in 185 limbs, 19 of which received reintervention. RFA also resulted in improved pain and less bruising compared to ligation and stripping in some studies (Hinchliffe, et al., 2006). Early studies, in addition to the more recent studies cited above, do support the safety and efficacy of RFA for the treatment of symptomatic saphenous varicosities, and is considered an appropriate alternative to conventional procedures.

ECRI Institute published an evidence report evaluating endovenous radio-frequency ablation (VNUS) for the treatment of varicose veins (ECRI, 2006). After reviewing the available evidence ECRI concluded that RFA offered a less invasive alternative to surgical stripping and ligation for patients with symptomatic varicose veins. ECRI noted that patients returned to work sooner and suffered less pain and fewer infections. Nonetheless, the benefits of RFA compared to surgery were supported on follow-up periods that were short term and consisted of a few days to one month posttreatment.

In 2003 NICE issued an Interventional Procedure Guidance for RFA and reported that safety and efficacy appeared adequate to support use of the procedure as an alternative to sapheno-femoral ligation and stripping.

Endovenous Laser Therapy (EVLT): EVLT, also commonly referred to as endovenous laser ablation of the saphenous vein (ELAS), is a treatment alternative to surgical stripping of the greater saphenous vein. EVLT is also considered an effective treatment for the SSV (Bhayani, Lippitz, 2009) however it is not typically used for smaller veins. EVLT is performed by threading a catheter through the greater saphenous vein and inserting an optical fiber through the catheter. The optical fiber is then connected to a surgical laser, allowing high-intensity laser light to induce photocoagulation of blood and occlusion of the vein. As the catheter is withdrawn, light pulses can be repeated at regular intervals to prevent any further blood flow through the vein. The procedure is typically used to treat larger varicose veins since catheters cannot be easily passed through a tortuous vein or a vein with several turns or bends. Small dilated branches that persist after EVLT may require additional treatments with sclerotherapy or phlebectomy (Radiological Society of North America, 2009).

The FDA has granted several approvals for ablative technologies, including: Diomed 810nm laser (Diomed, Inc.); Dornier diode laser systems (Dornier MedTech, Kennesaw, GA); Biolitec, Inc. (East Longmeadow, MA); Angiodynamics, Inc. and Vascular Solutions Inc. (Minneapolis, MN).

Evidence in the medical literature evaluating EVLT for the treatment of saphenous vein reflux consists of both retrospective and prospective case series, published reviews, and randomized controlled clinical trials (Huisman, et al., 2009; Nijsten, et al., 2009; Kalteis, et al., 2008; Darwood, et al., 2008; Desmyttrere, et al., 2007; Sharif, et al., 2007; Gibson, et al., 2007; Rasmussen, et al., 2007; Ravi, et al., 2006; Puggioni, et al., 2006; Min, et al., 2003; Ho, 2003; Chang and Chua, 2002; Proebstle, et al., 2002; Navarro and Min, 2001). There is a growing body of evidence to suggest that more minimally invasive techniques, which include both RFA and EVLT, are beneficial in the treatment of varicose veins when used alone (van den Bos, et al, 2009; Ravi et al., 2006; Sadick, 2005; Beale, et al., 2004; Teruya and Ballard, 2004; Elias and Frasier, 2004). Sample size and follow-up periods vary widely across studies; follow-up periods typically range at least one to four years on average. In some of the studies, duplex ultrasound demonstrated successful vein occlusion after initial treatment and throughout the various follow-up periods (Kalteis, et al., 2008; Gibson, et al., 2007; Desmyttrere, et al., 2007; Ravi, et al., 2006; Puggioni, et al., 2006; Min, et al., 2003). Some of the measured outcomes, such as complication rates, return to work, patient satisfaction and quality of scores, are mixed—some authors report improvement compared to traditional surgical methods while others have not. Success rates and recurrence rates have been promising with several studies supporting clinical efficacy. Van den Bos, et al. (2009) published the results of a meta-analysis demonstrating success rates of 78%, 84%, and 95% for ultrasound guided sclerotherapy, RFA and EVLT respectively, after three years. Min and associates (2003) reported a recurrence rate of less than 7% at a two-year follow-up, although the study had a significant number of patients lost to follow-up. Nonetheless, the authors noted their results were comparable or superior to those reported for other treatment options, including surgery, ultrasound-guided sclerotherapy, and radiofrequency ablation. Puggioni et al. (2006) concluded from a retrospective review that the overall success rate of endovenous ablation techniques for occluding the incompetent greater saphenous vein was 94% at one month, although the EVLT group developed more frequent postoperative complications compared to an RFA group. Ravi et al., (2006) reported that no GSV recanalization was found at three years post EVLT and that no saphenous vein could be identified in 82.5% of limbs in their study group. Closure rates at one month, one year, two year, three year and four year follow-up were reported by Desmyttrere, et al. (2007) as follows: 98.4%, 96.8%, 97.8%, 99.3% and 97.1%, respectively. Overall, much of the evidence available suggests that endovenous closure techniques are as good as or superior to conventional ligation and stripping of the greater saphenous vein.

ECRI Institute published an evidence report evaluating laser ablation of the greater saphenous vein (ECRI, 2004) and concluded that based on the available evidence endovenous laser ablation effectively occluded the greater saphenous vein for up to one year following treatment, complications were mild, and the retreatment rates were low. Data on quality of life was lacking.

NICE issued an Interventional Procedure Guidance for EVLT of the long saphenous vein. The guidance committee accepts the evidence on safety and efficacy as adequate to support the use of this procedure (NICE, 2004b). The evidence for efficacy was based on five case series with a mean follow-up of one to 17 months. Saphenous vein closure rates were between 90% and 100%. The authors noted that although procedure seems

effective in occluding the vein, few studies have reported on patient-oriented outcomes such as improvement in symptoms.

A position statement issued by the Society of Interventional Radiology in December 2003 calls the use of endovenous ablation therapy, performed with either laser or radiofrequency devices under imaging guidance and monitoring, an effective treatment of extremity venous reflux and varicose veins. The statement reports that the success rate for vein ablation ranges from 90–95% and that long-term results demonstrate recurrence rates of less than 7% at two-year follow-up. Lower rates of recurrence may be the result of the fact that imaging guidance enhances the ability to target and treat only the abnormal, incompetent venous segments. The society recommends using Duplex ultrasound prior to the procedure to map the necessary anatomy of the venous system, during the procedure for correct catheter placement and anesthetic delivery, and as necessary for follow-up. Currently, the 2003 position statement remains unchanged.

Endomechanical Ablative Approach: Minimally invasive treatment methods for treatment of varicose veins continue to evolve. One method under current investigation is the endomechanical ablative approach to varicose vein treatment utilizing a percutaneous infusion catheter. This approach involves the use of a special catheter (ClariVein™ [Vascular Insights, LLC, Madison, CT]) which combines two modalities of treatment for varicose veins: endovenous mechanical vein destruction with a rotating wire and the simultaneous infusion of an FDA approved liquid sclerosant, sodium tetradecyl sulfate to enhance venous occlusion. This mechanical-chemical ablative modality (endomechanical ablative approach) is described as minimally invasive and purported to accomplish great saphenous vein occlusion without the use of tumescent anesthesia. Information available from the manufacturer of ClariVein indicates the catheter has received 510(k) clearance from the United States Food and Drug Administration (FDA) for infusion of physician-specified agents in the peripheral vasculature. Currently evidence in the peer-reviewed published scientific literature supporting safety and efficacy is lacking, further studies are needed to support the clinical utility of this approach. .

Subfascial Endoscopic Perforator Surgery (SEPS): SEPS is a minimally invasive procedure for treating chronic venous insufficiency, in which incompetent perforating veins located in the calf are believed to be a contributing factor. Incompetent perforator veins result in pooling of blood in the lower extremity area, leading to vein enlargement, pain, swelling, skin discoloration and ulcers, and typically lead to chronic venous insufficiency.

An alternative to open subfascial perforator vein surgery (i.e., the Linton procedure), SEPS is recommended for patients in whom conservative measures have failed to treat chronic venous insufficiency and ulceration. The Linton procedure has been associated with a high incidence of postoperative wound healing complications (Townsend, 2004). Direct visualization through endoscopy has been suggested as a more desirable approach than the Linton technique. During SEPS, an endoscope is inserted in an incision located away from the ulcer site, and a balloon dissection is performed. The veins are ligated with clips and subsequently dissected, reducing pressure. Authors claim that stasis ulcer healing rates and maintenance of healing at five years after SEPS are 90% for patients with normally functioning deep venous systems and 75–80% for patients with deep venous insufficiencies (Elias, Frazier, 2004; Gloviczki, et al., 1999). The overall goal of SEPS in treating chronic venous ulcers is to interrupt the incompetent perforating veins in order to decrease reflux and pressure in areas above the ankle.

Evidence in the form of randomized clinical trials and both retrospective and prospective case series support the safety and efficacy of SEPS as an alternative to open procedures when performed for the treatment of incompetent medial calf perforator veins (Nelzen, Fransson, 2007; Kianifard, et al., 2007; de Rijcke, et al., 2003; Lee, et al., 2003; Kalra and Gloviczki, 2003; Sybrandy, et al., 2001; Pierek, et al., 1997). In contrast, SEPS performed for the treatment of post-thrombotic syndrome is controversial. Studies indicate that SEPS produces poorer outcomes, specifically, less ulcer healing and higher recurrence rates when used to treat limbs with post-thrombotic syndrome than when used to treat limbs with peripheral vascular insufficiency (Gloviczki, et al., 1999). -

NICE issued an Interventional Procedure Guidance for subfascial endoscopic perforator vein surgery. One randomized controlled trial, two non-randomized comparative studies and two case series were reviewed. The NICE specialty advisors noted that based on the evidence reviewed, efficacy of the procedure is unproven and the indications are not well established. Reported complications include nerve injury and deep vein thrombosis. There was evidence to support lower wound infection rates compared to the open procedure. Length of stay

was shorter for SEPS. The rate of primary ulcer healing and cumulative ulcer recurrence rates was comparable for both open and SEPS procedures. Although SEPS has been used for individuals with post-thrombotic valvular incompetence, there is evidence when used for this indication individuals may have poorer outcomes compared with individuals with primary valvular incompetence. In summary, the advisors noted careful patient selection is particularly important and there are uncertainties regarding safety of the procedure (NICE, 2004c).

Summary

The etiology of varicose veins is multifactorial and may result in a variety of symptoms and complications. Several treatment options are available, including minimally invasive surgical methods. The two main treatment options are surgery and sclerotherapy; however, there is little published data comparing their effectiveness. The peer-reviewed scientific literature supports safety and efficacy of most procedures, and most patients benefit from treatment, although recurrences have been reported in the literature. While varicose vein surgery is a very common surgical procedure, there is no general consensus regarding the best surgical approach.

Coding/Billing Information

Note: This list of codes may not be all-inclusive.

The treatment of varicose veins is covered only when coverage is available under the plan for varicose vein treatment. Benefit exclusions and limitations may apply. Invasive treatment of varicose veins is excluded under many plans and therefore the services listed below may not be covered.

The following are covered diagnoses when reported with a medically necessary and covered varicose vein procedure code:

ICD-9-CM Diagnosis Codes	Description
451.0	Phlebitis and thrombophlebitis of superficial vessels of lower extremities
451.11	Phlebitis and thrombophlebitis of femoral vein (deep) (superficial)
451.19	Phlebitis and thrombophlebitis of other deep vessels of lower extremities
453.40	Acute venous embolism and thrombosis of unspecified deep vessels of lower extremity
453.41	Acute venous embolism and thrombosis of deep vessels of proximal lower extremity
453.42	Acute venous embolism and thrombosis of deep vessels of distal lower extremity
453.50	Chronic venous embolism and thrombosis of unspecified deep vessels of lower extremity
453.51	Chronic venous embolism and thrombosis of unspecified deep vessels of proximal lower extremity
453.52	Chronic venous embolism and thrombosis of deep vessels of distal lower extremity
453.6	Venous embolism and thrombosis of superficial vessels of lower extremity
454.0-454.9	Varicose veins of lower extremities
459.11	Postphlebitic syndrome with ulcer
459.12	Postphlebitic syndrome with inflammation
459.13	Postphlebitic syndrome with ulcer and inflammation
459.19	Postphlebitic syndrome with other complication
459.2	Compression of vein
459.31	Chronic venous hypertension with ulcer
459.32	Chronic venous hypertension with inflammation
459.33	Chronic venous hypertension with ulcer and inflammation
682.6	Cellulitis and abscess of leg, except foot
682.7	Cellulitis of foot, except toes
707.10-	Ulcer of lower limb, except for pressure ulcer

707.19	
707.9	Chronic ulcer of unspecified site
729.5	Pain in limb
729.81	Swelling of limb
729.92	Nontraumatic hematoma of soft tissue
747.64	Lower limb vessel anomaly
782.3	Edema
891.1	Open wound of knee, leg [except thigh], and ankle, complicated
892.1	Open wound of foot, except toes, complicated
V12.52	Thrombophlebitis

Covered when medically necessary:

Sclerotherapy

CPT[®]* Codes	Description
36470	Injection of sclerosing solution; single vein
36471	Injection of sclerosing solution; multiple veins, same leg

HCPCS Codes	Description
S2202	Echosclerotherapy

Radiofrequency Ablation

CPT[®]* Codes	Description
36475	Endovenous ablation therapy of incompetent vein, extremity, inclusive of all imaging guidance and monitoring, percutaneous, radiofrequency; first vein treated
36476	Second and subsequent veins treated in a single extremity, each through separate access sites (List separately in addition to code for primary procedure)
37250	Intravascular ultrasound (non-coronary vessel) during diagnostic evaluation and/or therapeutic intervention; initial vessel (List separately in addition to code for primary procedure)
37251	Intravascular ultrasound (non-coronary vessel) during diagnostic evaluation and/or therapeutic intervention; each additional vessel (List separately in addition to code for primary procedure)
76942	Ultrasound guidance for needle placement (eg biopsy, aspiration, injection, localization device), imaging supervision and interpretation

Endovenous Laser Ablation

CPT[®]* Codes	Description
36478	Endovenous ablation therapy of incompetent vein, extremity, inclusive of all imaging guidance and monitoring, percutaneous, laser, first vein treated
36479	Second and subsequent veins treated in a single extremity, each through separate access sites (List separately in addition to code for primary procedure)
37250	Intravascular ultrasound (non-coronary vessel) during diagnostic evaluation and/or therapeutic intervention; initial vessel (List separately in addition to code for primary procedure)
37251	Intravascular ultrasound (non-coronary vessel) during diagnostic evaluation and/or therapeutic intervention; each additional vessel (List separately in addition to code for primary procedure)

76942	Ultrasound guidance for needle placement (eg biopsy, aspiration, injection, localization device), imaging supervision and interpretation
-------	--

Ligation and Excision

CPT^{®*} Codes	Description
37700	Ligation and division of long saphenous vein at saphenofemoral junction, or distal interruptions
37718	Ligation, division, and stripping, short saphenous vein
37722	Ligation, division, and stripping, long (greater) saphenous veins from saphenofemoral junction to knee or below
37735	Ligation and division and complete stripping of long or short saphenous veins with radical excision of ulcer and skin graft and/or interruption of communicating veins of lower leg, with excision of deep fascia

Ambulatory Phlebectomy

CPT^{®*} Codes	Description
37765	Stab phlebectomy of varicose veins, one extremity; 10-20 stab incisions
37766	Stab phlebectomy of varicose veins, one extremity; more than 20 incisions
37780	Ligation and division of short saphenous vein at saphenopopliteal junction (separate procedure)
37785	Ligation, division, and/or excision of varicose vein cluster(s), one leg
37799 [†]	Unlisted procedure, vascular surgery

[†]**Note:** Covered when medically necessary and used to report stab phlebectomy of varicose veins, one extremity; less than 10 incisions.

Subfascial Endoscopic Perforator Surgery

CPT* Codes	Description
37500	Vascular endoscopy, surgical, with ligation of perforator veins, subfascial (SEPS)
37760	Ligation of perforator veins, subfascial, radical (Linton type), with or without skin graft, open
37761	Ligation of perforator vein(s), subfascial, open, including ultrasound guidance, when performed, 1 leg

Not Covered/Not Medically Necessary/Cosmetic:

CPT* Codes	Description
36468	Single or multiple injections of sclerosing solutions, spider veins (telangiectasia); limb or trunk
36469	Single or multiple injections of sclerosing solutions, spider veins (telangiectasia); face

ICD-9-CM Diagnosis Codes	Description
	All other codes

Experimental/Investigational/Unproven/Not Covered when used to report endomechanical ablative approach for treatment of varicose veins:

CPT* Codes	Description
75894	Transcatheter therapy, embolization, any method, radiological supervision and interpretation
37204	Transcatheter occlusion or embolization (eg, for tumor destruction, to achieve hemostasis, to occlude a vascular malformation), percutaneous, any method, non-central nervous system, non-head or neck

ICD-9-CM Diagnosis Codes	Description
443.9	Peripheral vascular disease, unspecified
451.0	Phlebitis and thrombophlebitis of superficial vessels of lower extremities
451.11	Phlebitis and thrombophlebitis of femoral vein (deep) (superficial)
451.19	Phlebitis and thrombophlebitis of other deep vessels of lower extremities
451.9	Phlebitis and thrombophlebitis, of unspecified site
453.40	Acute venous embolism and thrombosis of unspecified deep vessels of lower extremity
453.41	Acute venous embolism and thrombosis of deep vessels of proximal lower extremity
453.42	Acute venous embolism and thrombosis of deep vessels of distal lower extremity
453.50	Chronic venous embolism and thrombosis of unspecified deep vessels of lower extremity
453.51	Chronic venous embolism and thrombosis of unspecified deep vessels of proximal lower extremity
453.52	Chronic venous embolism and thrombosis of deep vessels of distal lower extremity
453.6	Venous embolism and thrombosis of superficial vessels of lower extremity
453.89	Acute venous embolism and thrombosis of other specified veins
453.9	Acute venous embolism and thrombosis of unspecified site
454.0-454.9	Varicose veins of lower extremities
459.11	Postphlebitic syndrome with ulcer
459.12	Postphlebitic syndrome with inflammation
459.13	Postphlebitic syndrome with ulcer and inflammation
459.19	Postphlebitic syndrome with other complication
459.2	Compression of vein
459.31	Chronic venous hypertension with ulcer
459.32	Chronic venous hypertension with inflammation
459.33	Chronic venous hypertension with ulcer and inflammation
682.6	Cellulitis and abscess of leg, except foot
682.7	Cellulitis of foot, except toes
707.10-707.19	Ulcer of lower limb, except for pressure ulcer
707.9	Chronic ulcer of unspecified site
709.9	Unspecified disorder of skin and subcutaneous tissue
719.86	Other specified disorders of lower leg joint
729.5	Pain in limb
729.81	Swelling of limb
729.92	Nontraumatic hematoma of soft tissue
747.60	Anomaly of the peripheral vascular system, unspecified site
747.64	Lower limb vessel anomaly
782.3	Edema
891.1	Open wound of knee, leg [except thigh], and ankle, complicated
892.1	Open wound of foot, except toes, complicated
997.79	Vascular complications of other vessels
V12.52	Thrombophlebitis

References

1. Ahmad I, Ahmad W, Dingui M. Prevention or reversal of deep venous insufficiency by aggressive treatment of superficial venous disease. *Am J Surg*. 2006 Jan;191(1):33-8.
2. Alaiti S. Sclerotherapy: Treatment. *eMedicine Specialties*. April 16, 2010. Accessed October 13, 2010. Available at URL address: <http://emedicine.medscape.com/article/1271091-treatment>
3. Almeida JI, Kaufman J, Göckeritz O, Chopra P, Evans MT, Hoheim DF, et al. Radiofrequency endovenous ClosureFAST versus laser ablation for the treatment of great saphenous reflux: a multicenter, single-blinded, randomized study (RECOVERY study). *J Vasc Interv Radiol*. 2009 Jun;20(6):752-9.
4. Almeida JI, Raines JK. Radiofrequency ablation and laser ablation in the treatment of varicose veins. *Ann Vasc Surg*. 2006 Jul;20(4):547-52.
5. American Academy of Cosmetic Surgery. 2003 Guidelines for sclerotherapy. Accessed September 29, 2010. Available at URL address: <http://www.cosmeticsurgery.org/media/position.cfm>
6. American College of Phlebology. Guidelines for varicose vein surgery. 2008. Accessed October 12, 2010. Available at URL address: <http://vejledning.eu/filer%20til%20fag-ret/Varicose-Vein-Rx-Guidelines.pdf>
7. Angermeier MC. Treatment of facial vascular lesions with intense pulsed light. *J Cutan Laser Ther*. 1999 Apr;1(2):95-100.
8. Angle N, Freischlag JA. Venous disease. In: Townsend CM Jr., Beauchamp RD, Evers BM, Mattox KL, editors. *Sabiston textbook of surgery*. 17th ed. Philadelphia, PA: W.B. Saunders Co.; 2004. ch 67. Copyright © 2004 Elsevier.
9. Aremu MA, Mahendran B, Butcher W, Khan Z, Colgan MP, Moore DJ, et al. Prospective randomized controlled trial: conventional versus powered phlebectomy. *J Vasc Surg*. 2004 Jan;39(1):88-94.
10. Barrett JM, Allen B, Ockelford A, Goldman MP. Microfoam ultrasound-guided sclerotherapy of varicose veins in 100 legs. *Dermatol Surg*. 2004 Jan 1;30(1):6-12.
11. Beale RJ, Mavor AI, Gough MJ. Minimally invasive treatment for varicose veins: a review of endovenous laser treatment and radiofrequency ablation. *Int J Low Extrem Wounds*. 2004 Dec;3(4):188-97.
12. Belcaro G, Cesarone MR, Di Renzo A, Brandolini R, Coen L, Acerbi G, et al. Foam-sclerotherapy, surgery, sclerotherapy, and combined treatment for varicose veins: a 10-year, prospective, randomized, controlled trial (VEDICO trial). *Angiology*. 2003 May 1;54(3):307-15.
13. Bhayani R, Lippitz J. Varicose veins. *Dis Mon*. 2009 April; 55:212-22.
14. Bountouroglou DG, Azzam M, Kakkos SK, Pathmarajah M, Young P, Geroulakos G. Ultrasound-guided foam sclerotherapy combined with sapheno-femoral ligation compared to surgical treatment of varicose veins: early results of a randomized controlled trial. *Eur J Vasc Endovasc Surg*. 2006 Jan;31(1):93-100.
15. Bowes LE, Goldman MP. Sclerotherapy of reticular and telangiectatic veins of the face, hands, and chest. *Dermatol Surg*. 2002 Jan;28(1):46-51.
16. Chandler JG, Pichot O, Sessa C, Schuller-Petrovic, Kavnick L, Betgan J. Treatment of primary venous insufficiency by endovenous saphenous vein obliteration. *Vasc Surg*. 2000;34:201-14.

17. Chang CJ, Chua JJ. Endovenous laser photocoagulation (EVLP) for varicose veins. *Lasers Surg Med.* 2002;31(4):257-62.
18. Cheshire N, Elias SM, Keagy B, Kolvenbach R, Leahy AL, Marston W, Pannier-Fischer F, Rabe E, Spitz GA. Powered phlebectomy (TriVex) in treatment of varicose veins. *Ann Vasc Surg.* 2002 Jul;16(4):488-94.
19. Chetter IC, Mylankal KJ, Hughes H, Fitridge R. Randomized clinical trial comparing multiple stab incision phlebectomy and transilluminated powered phlebectomy for varicose veins. *Br J Surg.* 2006 Feb;93(2):169-74.
20. Corabian P, Harstall C. Sclerotherapy for leg varicose veins. Alberta Heritage Foundation for Medical Research. Technology Assessment. Ip-19 Information paper. May 2004. ©Copyright Alberta Heritage Foundation for Medical Research, 2004.
21. Creton D, Uhl JF. Foam sclerotherapy combined with surgical treatment for recurrent varicose veins: short term results. *Eur J Vasc Endovasc Surg.* 2007 May;33(5):619-24.
22. Darke SG, Baker SJ. Ultrasound-guided foam sclerotherapy for the treatment of varicose veins. *Br J Surg.* 2006 Aug;93(8):969-74.
23. Darvall KA, Bate GR, Silverman SH, Adam DJ, Bradbury AW. Medium-term results of ultrasound-guided foam sclerotherapy for small saphenous varicose veins. *Br J Surg.* 2009 Nov;96(11):1268-73.
24. Darwood RJ, Theivacumar N, Dellagrammaticas D, Mavor AI, Gough MJ. Randomized clinical trial comparing endovenous laser ablation with surgery for the treatment of primary great saphenous varicose veins. *Br J Surg.* 2008 Mar;95(3):294-301.
25. de Rijcke PA, Hop WC, Wittens CH. Subfascial endoscopic perforating vein surgery as treatment for lateral perforating vein incompetence and venous ulceration. *J Vasc Surg.* 2003 Oct;38(4):799-803.
26. Desmytère J, Grard C, Wassmer B, Mordon S. Endovenous 980-nm laser treatment of saphenous veins in a series of 500 patients. *J Vasc Surg.* 2007 Dec;46(6):1242-7.
27. Disselhoff BC, der Kinderen DJ, Kelder JC, Moll FL. Randomized clinical trial comparing endovenous laser ablation of the great Saphenous vein with and without ligation of the sapheno-femoral junction: 2-year results. *Eur J Vasc Endovasc Surg.* 2008 Dec;36(6):713-8.
28. Disselhoff BC, der Kinderen DJ, Kelder JC, Moll FL. Randomized clinical trial comparing endovenous laser with cryostripping for great saphenous varicose veins. *Br J Surg.* 2008 Oct;95(10):1232-8.
29. Duffy DM. Sclerosants: a comparative review. *Dermatol Surg.* 2010 Jun;36 Suppl 2:1010-25.
30. Duffy DM, Garcia C, Clarke RE. The role of sclerotherapy in abnormal varicose hand veins. *Plast Reconstr Surg.* 1999 Oct;104(5):1474-9; discussion 1480-1.
31. ECRI Institute. Hotline Response [database online]. Plymouth Meeting (PA): ECRI Institute; Endovenous radiofrequency ablation (VNUS Closure System) for the treatment of varicose veins. 2008, Aug 8. Available at URL address: <http://www.ecri.org>.
32. ECRI Institute. Endovenous Radio-frequency Ablation (VNUS Closure System) for the Treatment of Varicose Veins. Plymouth Meeting (PA): ECRI Institute; 2006 August 24. (Evidence Report Issue No. 138). Available at URL address: <http://www.ecri.org>.
33. ECRI Institute. Endovenous Laser Ablation of the Greater Saphenous Vein. Plymouth Meeting (PA): ECRI Institute; 2004, December 30. (Evidence Report Issue No. 121). Available at URL address: <http://www.ecri.org>.

34. ECRI Institute. Transilluminated powered phlebectomy (TIPP) for varicose veins [Emerging Technology Evidence Report]. Plymouth Meeting (PA): ECRI Institute; 2008, April 8. Available at URL address: <http://www.ecri.org>.
35. Eklöf B, Rutherford RB, Bergan JJ, Carpentier PH, Gloviczki P, Kistner RL, et al., American Venous Forum International Ad Hoc Committee for Revision of the CEAP Classification. Revision of the CEAP classification for chronic venous disorders: consensus statement. *J Vasc Surg*. 2004 Dec;40(6):1248-52.
36. Feied C. Varicose vein treatment and endovenous laser therapy. *eMedicine specialties. Dermatology, Diseases of the vessels*. Updated July 7, 2010. Accessed September 29, 2010. Available at URL address: <http://www.emedicine.com/med/topic2788.htm>
37. Franz RW, Knapp ED. Transilluminated Powered Phlebectomy Surgery for Varicose Veins: A Review of 339 Consecutive Patients. *Ann Vasc Surg*. 2008 Sep 5. [Epub ahead of print.
38. Gibson KD, Ferris BL, Pepper D. Endovenous laser treatment of varicose veins. *Surg Clin North Am*. 2007a Oct;87(5):1253-65, xii.
39. Gibson KD, Ferris BL, Pepper D. Foam sclerotherapy for the treatment of superficial venous insufficiency. *Surg Clin North Am*. 2007b Oct;87(5):1285-95, xii-xiii.
40. Gibson KD, Ferris BL, Polissar N, Neradilek B, Pepper D. Endovenous laser treatment of the small [corrected] saphenous vein: efficacy and complications. *J Vasc Surg*. 2007 Apr;45(4):795-801; discussion 801-3.
41. Gloviczki P, Bergan JJ, Rhodes JM, Canton LG, Harmsen S, Ilstrup DM. Mid-term results of endoscopic perforator vein interruption for chronic venous insufficiency: lessons learned from the North American subfascial endoscopic perforator surgery registry. The North American Study Group. *J Vasc Surg*. 1999;29(3):489-502.
42. Goldman MP. Closure of the greater saphenous vein with endoluminal radiofrequency thermal heating of the vein wall in combination with ambulatory phlebectomy: preliminary 6-month follow-up. *Dermatol Surg*. 2000;26:452-56.
43. Gonzalez-Zeh R, Armisen R, Barahona S. Endovenous laser and echo-guided foam ablation in great saphenous vein reflux: one-year follow-up results. *J Vasc Surg*. 2008 Oct;48(4):940-6.
44. Habif: *Clinical Dermatology*, 5th ed, Ch 3. Stasis dermatitis and venous ulceration: postphlebotic syndromes. Copyright © 2009 Mosby.
45. Hamper UM, DeJong MR, Scoutt LM. Ultrasound Evaluation of the Lower Extremity Veins. *Radiol Clin North Am*. 2007 May;45(3):525-47.
46. Hinchliffe RJ, Ubhi J, Beech A, Ellison J, Braithwaite BD. A prospective randomized controlled trial of VNUS closure versus surgery for the treatment of recurrent long saphenous varicose veins. *Eur J Vasc Endovasc Surg*. 2006 Feb;31(2):212-8.
47. Ho P, Poon JT, Cho SY, Cheung G, Tam YF, Yuen WK, Cheng SW. Day surgery varicose vein treatment using endovenous laser. *Hong Kong Med J*. 2009 Feb;15(1):39-43.
48. Huisman LC, Bruins RM, van den Berg M, Hissink RJ. Endovenous laser ablation of the small saphenous vein: prospective analysis of 150 patients, a cohort study. *Eur J Vasc Endovasc Surg*. 2009 Aug;38(2):199-202.
49. Jia X, Mowatt G, Burr JM, Cassar K, Cook J, Fraser C. Systematic review of foam sclerotherapy for varicose veins. *Br J Surg*. 2007 Aug;94(8):925-36.

50. Kahle B, Leng K. Efficacy of sclerotherapy in varicose veins—prospective, blinded, placebo-controlled study. *Dermatol Surg*. 2004 May;30(5):723-8; discussion 728.
51. Kakkos SK, Bountouroglou DG, Azzam M, Kalodiki E, Daskalopoulos M, Geroulakos G. Effectiveness and safety of ultrasound-guided foam sclerotherapy for recurrent varicose veins: immediate results. *J Endovasc Ther*. 2006 Jun;13(3):357-64.
52. Kalra M, Gloviczki P. Surgical treatment of venous ulcers: role of subfascial endoscopic perforator vein ligation. *Surg Clin North Am*. Jun 2003;83(3):671-705.
53. Kalteis M, Berger I, Messie-Werndl S, Pistrich R, Schimetta W, Pölz W, Hieller F. High ligation combined with stripping and endovenous laser ablation of the great saphenous vein: early results of a randomized controlled study. *J Vasc Surg*. 2008 Apr;47(4):822-9; discussion 829.
54. Kendler M, Wetzig T, Simon JC. Foam sclerotherapy—a possible option in therapy of varicose veins. *J Dtsch Dermatol Ges*. 2007 Aug;5(8):648-54.
55. Kianifard B, Holdstock J, Allen C, Smith C, Price B, Whiteley MS. Randomized clinical trial of the effect of adding subfascial endoscopic perforator surgery to standard great saphenous vein stripping. *Br J Surg*. 2007 Sep;94(9):1075-80.
56. Klem TM, Schnater JM, Schütte PR, Hop W, van der Ham AC, Wittens CH.. A randomized trial of cryo stripping versus conventional stripping of the great saphenous vein. *J Vasc Surg*. 2009 Feb;49(2):403-9.
57. Kouri B. Current evaluation and treatment of lower extremity varicose veins. *Am J Med*. 2009 Jun;122(6):513-5.
58. Labas P, Ohradka B, Cambal M, Reis R, Fillo J. Long term results of compression sclerotherapy. *Bratisl Lek Listy*. 2003;104(2):78-81.
59. Leach BC, Goldman MP. Comparative trial between sodium tetradecyl sulfate and glycerin in the treatment of telangiectatic leg veins. *Dermatol Surg*. 2003 Jun;29(6):612-4; discussion 615.
60. Lee BJ. The role of sclerotherapy in abnormal varicose hand veins. *Plast Reconstr Surg*. 2000 Jul;106(1):227-9.
61. Lee DW, Lam YH, Chan AC, Chung SC. Subfascial endoscopic perforator surgery for venous ulcers. *Hong Kong Med J*. 2003 Aug;9(4):279-82.
62. Leopardi D, Hoggan BL, Fitridge RA, Woodruff PW, Maddern GJ. Systematic review of treatments for varicose veins. *Ann Vasc Surg*. 2009 Mar;23(2):264-76.
63. Lew WK, Weaver FA. Varicose Veins: Treatment. *eMedicine Specialties*. Updated July 7, 2010. Accessed October 13, 2010. Available at URL address: <http://emedicine.medscape.com/article/462579-treatment>
64. Luebke T, Brunkwall J. Meta-analysis of transilluminated powered phlebectomy for superficial varicosities. *J Cardiovasc Surg (Torino)*. 2008 Dec;49(6):757-64.
65. Lurie F, Cretan D, Ekloff B, Kabnick LS, Kistner RI, Pichot O, et al. Prospective randomized study of endovenous radiofrequency obliteration (closure procedure) versus ligation and stripping in selected patient population (EVOLVeS Study). *J Vasc Surg*. 2003;Aug 38(2):207-14.
66. Lurie F, Cretan D, Ekloff B, Kabnick LS, Kistner RL, Pichot O, et al. Prospective randomised study of endovenous radiofrequency obliteration (Closure) versus ligation and vein stripping (EVOLVeS): Two-year follow-up. *Eur J Vasc Endovasc Surg*. 2005 Jan;29:67-73.

67. McDonagh B, Sorenson S, Gray C, Huntley DE, Putterman P, King T, et al. Clinical spectrum of recurrent postoperative varicose veins and efficacy of sclerotherapy management using the compass technique. *Phlebology*. 2003;18:173-86.
68. Menyhei G, Gyevnár Z, Arató E, Kelemen O, Kollár L. Conventional stripping versus cryostripping: a prospective randomised trial to compare improvement in quality of life and complications. *Eur J Vasc Endovasc Surg*. 2008 Feb;35(2):218-23.
69. Merchant RF, Pichot O, for the Closure Study Group. Long-term outcomes of endovenous radiofrequency obliteration of saphenous reflux as a treatment for superficial venous insufficiency. *J Vasc Surg*. 2005 Sep;42(3):502-9, discussion 509.
70. Michaels JA, Campbell WB, Brazier JE, Macintyre JB, Ratcliffe J, Rigby K. Randomised clinical trial, observational study and assessment of cost-effectiveness of the treatment of varicose veins (REACTIV trial). *Health Technol Assess*. 2006 Apr;10(13):1-196, iii-iv.
71. Min RJ, Khilnani N, Zimmet SE. Endovenous laser treatment of saphenous vein reflux: long term results. *J Vasc Interv Radiol*. 2003 Aug;14(8):991-6.
72. Min RJ, Navarro L. Transcatheter duplex ultrasound-guided sclerotherapy for treatment of greater saphenous vein reflux: preliminary report. *Dermatol Surg*. 2000 May;26(5):410-4; discussion 413-4.
73. Murray JD, Bergan JJ, Riffenburgh RH. Development of open-scope subfascial perforating vein surgery: lessons learned from the first 67 cases. *Ann Vasc Surg*. 1999 Jul;13(4):372-7.
74. National Institute for Clinical Excellence (NICE). Endovenous laser treatment of the long saphenous vein. Guidance. Issued 2004b Mar 4. Accessed September 29, 2010. Available at URL address: <http://www.nice.org.uk/search/searchresults.jsp?keywords=varicose+vein&searchType=all>
75. National Institute for Clinical Excellence (NICE). Radiofrequency ablation of varicose veins. Guidance. Issued September 2003. Accessed September 29, 2010. Available at URL address: <http://www.nice.org.uk/search/searchresults.jsp?keywords=varicose+vein&searchType=all>
76. National Institute for Clinical Excellence (NICE). Subfascial endoscopic perforator surgery, Guidance. Issued 2004c June. Accessed September 29, 2010. Available at URL address: <http://www.nice.org.uk/search/searchresults.jsp?keywords=varicose+vein&searchType=all>
77. National Institute for Clinical Excellence (NICE). Transilluminated powered phlebectomy for varicose veins, Guidance. Issued 2004a Jan. Accessed September 29, 2010. Available at URL address: <http://www.nice.org.uk/search/searchresults.jsp?keywords=varicose+vein&searchType=all>
78. National Institute for Clinical Excellence (NICE). Ultrasound guided foam sclerotherapy for varicose veins. Guidance. Issued June 2006. Re-issued May 2007. Accessed September 29, 2010. Available at URL address: <http://www.nice.org.uk/search/searchresults.jsp?keywords=varicose+vein&searchType=all>
79. Navarro L, Min RJ, Bon E. Endovenous laser: a new minimally invasive method of treatment for varicose veins -- preliminary observations using an 810 nm Diode Laser. *Dermatol Surg*. 2001; 27(2):117-22.
80. Nelzén O, Fransson I. True long-term healing and recurrence of venous leg ulcers following SEPS combined with superficial venous surgery: a prospective study. *Eur J Vasc Endovasc Surg*. 2007 Nov;34(5):605-12.
81. Nijsten T, van den Bos RR, Goldman MP, Kockaert MA, Proebstle TM, Rabe E, et al. Minimally invasive techniques in the treatment of saphenous varicose veins. *J Am Acad Dermatol*. 2009 Jan;60(1):110-9.

82. Ogawa T, Hoshino S, Midorikawa H, Sato K. Clinical results of radiofrequency endovenous obliteration for varicose veins. *Surg Today*. 2005;35(1):47-51.
83. Ozkan U. Endovenous laser ablation of incompetent perforator veins: a new technique in treatment of chronic venous disease. *Cardiovasc Intervent Radiol*. 2009 Sep;32(5):1067-70.
84. Passman MA, Dattilo JB, Guzman RJ, Naslund TC. Combined endovenous ablation and transilluminated powered phlebectomy: is less invasive better? *Vasc Endovascular Surg*. 2007 Feb-Mar;41(1):41-7.
85. Peden E, Lumsden A. Radiofrequency ablation of incompetent perforator veins. *Perspect Vasc Surg Endovasc Ther*. 2007 Mar;19(1):73-7.
86. Perkowski P, Ravi R, Gowda RC, Olsen D, Ramaiah V, Rodriguez-Lopez JA, Diethrich EB. Endovenous laser ablation of the saphenous vein for treatment of venous insufficiency and varicose veins: early results from a large single-center experience. *Endovasc Ther*. 2004 Apr;11(2):132-8.
87. Pierik EG, van Urk H, Wittens CH. Efficacy of subfascial endoscopy in eradicating perforating veins of the lower leg and its relation with venous ulcer healing. *J Vasc Surg*. 1997 Aug;26(2):255-9.
88. Proebstle TM, Herdemann S. Early results and feasibility of incompetent perforator vein ablation by endovenous laser treatment. *Dermatol Surg*. 2007 Feb;33(2):162-8.
89. Proebstle TM, Lehr HA, Kargl A, Espinola-Klein C, Rother W, Bethge S, Knop J. Endovenous treatment of the greater saphenous vein with 940-nm diode laser: thrombotic occlusion after endoluminal thermal damage by laser-generated steam bubbles. *J Vasc Surg*. 2002;35(4):729-36.
90. Puggioni A, Kalra M, Carmo M, Mozes G, Gloviczki P. Endovenous laser therapy and radiofrequency ablation of the great saphenous vein: analysis of early efficacy and complications. *J Vasc Surg*. 2005 Sep;42(3):488-93.
91. Rabe E, Pannier F. Sclerotherapy of varicose veins with polidocanol based on the guidelines of the German Society of Phlebology. *Dermatol Surg*. 2010 Jun;36 Suppl 2:968-75.
92. Rabe E, Pannier-Fischer F, Gerlach H, Breu FX, Guggenbichler S, Zabel M; German Society of Phlebology. Guidelines for sclerotherapy of varicose veins (ICD 10: I83.0, I83.1, I83.2, and I83.9). *Dermatol Surg*. 2004 May;30(5):687-93; discussion 693.
93. Rabe E, Otto J, Schliephake D, Pannier F. Efficacy and safety of great saphenous vein sclerotherapy using standardised polidocanol foam (ESAF): a randomised controlled multicentre clinical trial. *Eur J Vasc Endovasc Surg*. 2008 Feb;35(2):238-45.
94. Radiological Society of North America. Varicose vein treatment (Endovenous ablation of varicose vein). June 29, 2009. Reviewed February 10, 2010. Copyright © 2010 Radiological Society of North America, Inc. (RSNA). Accessed September 29, 2010. Available at URL address: <http://www.radiologyinfo.org/en/info.cfm?PG=varicoseabl>
95. Rao J, Wildemore JK, Goldman MP. Double-blind prospective comparative trial between foamed and liquid polidocanol and sodium tetradecyl sulfate in the treatment of varicose and telangiectatic leg veins. *Dermatol Surg*. 2005 Jun;31(6):631-5; discussion 635.
96. Rasmussen LH, Bjoern L, Lawaetz M, Blemings A, Lawaetz B, Eklof B. Randomized trial comparing endovenous laser ablation of the great saphenous vein with high ligation and stripping in patients with varicose veins: short-term results. *J Vasc Surg*. 2007 Aug;46(2):308-15.
97. Raulin C, Weiss RA, Schonemark MP. Treatment of essential telangiectasias with an intense pulsed light source (PhotoDerm VL). *Dermatol Surg*. 1997 Oct;23(10):941-5.

98. Rautio R, Saarinen J, Laranne J, Salenius JP, Keski-Nisula L. Endovascular treatment of venous malformations in extremities: results of sclerotherapy and the quality of life after treatment. *Acta Radiol*. 2004 Jul;45(4):397-403.
99. Ravi R, Rodriguez-Lopez JA, Trayler EA, Barrett DA, Ramaiah V, Diethrich EB. Endovenous ablation of incompetent saphenous veins: a large single-center experience. *J Endovasc Ther*. 2006 Apr;13(2):244-8.
100. Rigby KA, Palfreyman SJ, Beverley C, Michaels JA. Surgery for varicose veins: use of tourniquet. *Cochrane Database of Systematic Reviews* 2002. In: *The Cochrane Library*, 2007, Issue 4. Copyright © 2007 The Cochrane Collaboration.
101. Rigby KA, Palfreyman SJ, Beverley C, Michaels JA. Surgery versus sclerotherapy for the treatment of varicose veins. *Cochrane Database of Systematic Reviews* 2004. In: *The Cochrane Library*, 2007 Issue 4. Copyright © 2007 The Cochrane Collaboration.
102. Roth SM. Endovenous radiofrequency ablation of superficial and perforator veins. *Surg Clin North Am*. 2007 Oct;87-5: 1267-84, xii.
103. Sadick NS. Advances in the treatment of varicose veins: Ambulatory phlebectomy, foam sclerotherapy, endovascular laser, and radiofrequency closure. *Dermatol Clin*. 2005 Jul;23(3):443-55,vi.
104. Scavee V. Transilluminated powered phlebectomy: not enough advantages? Review of the literature. *Eur J Vasc Endovasc Surg*. 2006 Mar;31(3):316-9. Epub 2005 Dec 15.
105. Scavee V, Lemaire E, Haxhe JP. Transilluminated powered phlebectomy. Mid-term clinical experience. *Int Angiol*. 2005 Mar;24(1):75-9.
106. Scavee V, Lesceu O, Theys S, Jamart J, Louagie Y, Schoevaerdt JC. Hook phlebectomy versus transilluminated powered phlebectomy for varicose vein surgery: early results. *Eur J Vasc Endovasc Surg*. 2003 May;25(5):473-5.
107. Shamiyeh A, Schrenk P, Huber E, Danis J, Wayand WU. Transilluminated powered phlebectomy: advantages and disadvantages of a new technique. *Dermatol Surg*. 2003 Jun;29(6):616-9.
108. Sharif MA, Lau LL, Lee B, Hannon RJ, Soong CV. Role of endovenous laser treatment in the management of chronic venous insufficiency. *Ann Vasc Surg*. 2007 Sep;21(5):551-5.
109. Singh MJ, Sura C. Endovenous saphenous and perforator vein ablation. Operative techniques in general Surgery. 2008 Sep; 10(3):131-135.
110. Society for Interventional Radiology. Position statement. Endovenous ablation. 2003 Dec. Accessed September 29, 2010. Available at URL address: http://www.sirweb.org/clinical/cpg/SIR_venous_ablation_statement_final_Dec03.pdf
111. Spitz GA, Braxton JM, Bergan JJ. Outpatient varicose vein surgery with transilluminated powered phlebectomy. *Vasc Surg*. 2000;34(6):547-55.
112. Subramonia S, Lees TA. The treatment of varicose veins. *Ann R Coll Surg Engl*. 2007 Mar;89(2):96-100.
113. Sybrandy JE, van Gent WB, Pierik EG, Wittens CH. Endoscopic versus open subfascial division of incompetent perforating veins in the treatment of venous leg ulceration: long-term follow-up. *J Vasc Surg*. 2001 May;33(5):1028-32.
114. Teruya TH, Ballard JL. New approaches for the treatment of varicose veins. *Surg Clin N Am*. 2004 Oct;84(5):1397-417viii-ix.

115. Tisi PV, Beverley CA. Injection sclerotherapy for varicose veins. Cochrane database of systematic reviews. In: The Cochrane Library, Issue 2, 2005. ©2005 The Cochrane Collaboration.
116. Trelles MA, Allones I, Alvarez J, Velez M, Martin-Vazquez M, Trelles OR, et al. The 800-nm diode laser in the treatment of leg veins: Assessment at 6 months. *J Am Acad Dermatol*. 2006 Feb;54(2):282-9.
117. Uchino IJ. Endovenous laser closure of the perforating vein of the leg. *Phlebology*. 2007;22(2):80-2.
118. Uurto I, Hannukainen J, Aarnio P. Single-center experience with foam sclerotherapy without ultrasound guidance for treatment of varicose veins. *Dermatol Surg*. 2007 Nov;33(11):1334-9; discussion 1339.
119. van den Bos R, Arends L, Kockaert M, Neumann M, Nijsten T. Endovenous therapies of lower extremity varicosities: a meta-analysis. *J Vasc Surg*. 2009 Jan;49(1):230-9.
120. VNUS Medical Technologies. Safety summary. Accessed September 29, 2010. Copyright © 2010 VNUS Medical Technologies. Available at URL address: <http://www.vnus.com/navigation/safety.htm>
121. Weiss RA. American College of Phlebology (ACP). Radio-frequency endovenous occlusion (Closure® technique). Accessed October 2003. Available at URL address: <http://www.phlebology.org/closure.htm>
122. Weiss RA, Dover JS. Laser surgery of leg veins. *Dermatol Clin*. 2002 Jan;20(1):19-36.
123. Weiss RA. Evaluation of the venous system by Doppler ultrasound and photoplethysmography or light reflection rheography before sclerotherapy. *Semin Dermatol*. 1993;12:78-87.
124. Weiss RA, Goldman MP. Transillumination mapping prior to ambulatory phlebectomy. *Dermatol Surg*. 1998 Apr;24(4):447-50.
125. Weiss RA, Heagle CR, Raymond-Martimbeau P. The Bulletin of the North American Society of Phlebology. Insurance Advisory Committee Report. *J Dermatol Surg Oncol*. 1992 Jul;18(7):609-16.
126. Weiss RA, Weiss MA. Controlled radiofrequency endovenous occlusion using a unique radiofrequency catheter under duplex guidance to eliminate saphenous varicose vein reflux: a 2-year follow-up. *Dermatol Surg*. 2002 Jan;28(1):38-42.
127. Welch HJ. Endovenous ablation of the great saphenous vein may avert phlebectomy for branch varicose veins. *J Vasc Surg*. 2006 Sep;44(3):601-5.
128. Welch HJ, Villavicencio JL. Primary varicose veins of the upper extremity: a report of three cases. *J Vasc Surg*. 1994 Nov;20(5):839-43.
129. Wright D, Gobin JP, Bradbury AW, Coleridge-Smith P, Spoelstra H, Berridge D, et al. Varisolve® polidocanol microfoam compared with surgery or sclerotherapy in the management of varicose veins in the presence of trunk vein incompetence: European randomized controlled trial. *Phlebology* 2006;21(4):180-190.

Policy History

Pre-Merger Organizations	Last Review Date	Policy Number	Title
CIGNA HealthCare	11/15/2007	0234	Varicose Vein Treatments
Great-West Healthcare	1/1/2007	04.202.03	Varicose Vein Treatment
	1/1/2007	04.258.02	Varicose Vein Treatment, Sclerotherapy

“CIGNA”, “CIGNA HealthCare” and the “Tree of Life” logo are registered service marks of CIGNA Intellectual Property, Inc., licensed for use by CIGNA Corporation and its operating subsidiaries. All products and services are provided by such operating subsidiaries and not by CIGNA Corporation. Such operating subsidiaries include Connecticut General Life Insurance Company, CIGNA Health and Life Insurance Company, CIGNA Behavioral Health, Inc., CIGNA Health Management, Inc., and HMO or service company subsidiaries of CIGNA Health Corporation and CIGNA Dental Health, Inc. In Arizona, HMO plans are offered by CIGNA HealthCare of Arizona, Inc. In California, HMO plans are offered by CIGNA HealthCare of California, Inc. In Connecticut, HMO plans are offered by CIGNA HealthCare of Connecticut, Inc. In North Carolina, HMO plans are offered by CIGNA HealthCare of North Carolina, Inc. In Virginia, HMO plans are offered by CIGNA HealthCare Mid-Atlantic, Inc. All other medical plans in these states are insured or administered by Connecticut General Life Insurance Company or CIGNA Health and Life Insurance Company.