

ACITRETIN

Products Affected

- Acitretin

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For initial therapy in the treatment of psoriasis: trial and failure, contraindication, or intolerance to methotrexate or cyclosporine is required. For continuation of therapy, approve if patient has already been started on Acitretin.

ACTIMMUNE

Products Affected

- Actimmune

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

ALIMTA

Products Affected

- Alimta

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	B vs D coverage determination

ALOSETRON

Products Affected

- Alosetron Hydrochloride

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Alosetron will not be approved for use in men, as safety and efficacy in men has not been established.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Alosetron is considered medically necessary for the treatment of severe IBS-D. At least one of the following must be present for diarrhea to be considered severe: frequent and severe abdominal pain or discomfort, frequent bowel urgency or fecal incontinence, and disability or restriction of daily activities due to IBS.

ANABOLIC STEROIDS, ANDROGENS

Products Affected

- Anadrol-50
- Oxandrolone TABS
- Testosterone GEL 25MG/2.5GM, 50MG/5GM
- Testosterone Pump GEL 1%

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

ANTIFUNGALS, AZOLE

Products Affected

- Voriconazole INJ
- Voriconazole SUSR
- Voriconazole TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documented fungal culture and/or notes from medical record suggestive of a serious fungal infection. For prophylactic use, fungal culture and medical records are not required.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 to 6 months, depending on indication
Other Criteria	For the treatment of oropharyngeal candidiasis, the candidiasis must be refractory to itraconazole or fluconazole.

ANTIFUNGALS, POLYENE

Products Affected

- Abelcet
- Ambisome
- Amphotericin B INJ

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	B vs D coverage determination

ANTIFUNGALS, SUPERFICIAL AND SYSTEMIC

Products Affected

- Caspofungin Acetate
- Itraconazole CAPS
- Itraconazole SOLN

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Onychomycosis (fingernails)-2mo. Onychomycosis (toenails)-3mo. All other indications -12 months
Other Criteria	For the treatment of tinea versicolor or pityriasis, use of oral ketoconazole or a topical antifungal agent is required prior to the use of Itraconazole. For candidiasis infections (unless specified <i>C. glabrata</i> or <i>C. krusei</i>), use of fluconazole is required prior to the use of itraconazole.

ANTIFUNGALS, TRIAZOLE

Products Affected

- Noxafil SUSP
- Noxafil TBEC
- Posaconazole Dr

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	<p>For the prophylaxis of invasive Aspergillus and Candida infections: Noxafil (posaconazole) is considered medically necessary in patients who are at high risk of developing these infections due to being severely immunocompromised, such as hematopoietic stem cell transplant (HSCT) recipients with graft-versus-host disease (GVHD) or those with hematologic malignancies with prolonged neutropenia from chemotherapy.</p> <p>For the treatment of oropharyngeal candidiasis, the candidiasis must be refractory to itraconazole or fluconazole.</p>

ANTINEOPLASTICS, MONOCLONAL ANTIBODIES

Products Affected

- Abraxane
- Aliqopa
- Avastin
- Bavencio
- Besponsa
- Bortezomib
- Cyramza
- Darzalex
- Gazyva
- Enhertu
- Herceptin
- Herceptin Hylecta
- Imfinzi
- Kadcyla
- Kanjinti INJ 420MG
- Keytruda INJ 100MG/4ML
- Lartruvo
- Libtayo
- Lumoxiti
- Mvasi
- Mylotarg
- Opdivo
- Perjeta
- Poteligeo
- Rituxan
- Rituxan Hycela
- Tecentriq
- Unituxin
- Vectibix INJ 100MG/5ML,
400MG/20ML
- Velcade
- Yervoy
- Yondelis

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	B vs D coverage determination

APOKYN

Products Affected

- Apokyn INJ 30MG/3ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

ARCALYST

Products Affected

- Arcalyst

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	12 years of age and older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	B vs D coverage determination

ARIKAYCE

Products Affected

- Arikayce

PA Criteria	Criteria Details
Indications	All FDA-Approved Indications, Some Medically-Accepted Indications.
Off-Label Uses	Cystic fibrosis pseudomonas aeruginosa infection
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous medication history
Age Restrictions	MAC-18 years and older
Prescriber Restrictions	MAC-Prescribed by a pulmonologist, infectious disease physician or a physician who specializes in the treatment of MAC lung infections. Cystic fibrosis-prescribed by or in consultation with a pulmonologist or physician who specializes in the treatment of cystic fibrosis
Coverage Duration	1 year
Other Criteria	MAC Lung disease-approve if the patient has NOT achieved negative sputum cultures for Mycobacterium avium complex after a background multidrug regimen AND Arikayce will be used in conjunction to a background multidrug regimen. Note-a multidrug regimen typically includes a macrolide (azithromycin or clarithromycin), ethambutol and a rifamycin (rifampin or rifabutin). Cystic fibrosis-patient has pseudomonas aeruginosa in culture of the airway.

AURYXIA

Products Affected

- Auryxia

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Diagnosis Iron deficiency anemia not on dialysis
Required Medical Information	Documentation of dialysis and hyperphosphatemia
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

AUSTEDO

Products Affected

- Austedo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Chorea associated with Huntington's disease: trial and failure, contraindication, or intolerance to tetrabenazine. For the treatment of tardive dyskinesia, no trial and failure is required

BANZEL

Products Affected

- Banzel

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

BENLYSTA

Products Affected

- Benlysta INJ 120MG, 400MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Concurrent use with other biologics or with cyclophosphamide intravenous (IV)
Required Medical Information	The patient must have a positive autoantibody test (i.e., anti-nuclear antibody [ANA] greater than or equal to 1:80 and/or anti-double-stranded DNA [anti-dsDNA] greater than or equal to 30 IU/ml).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	The patient must be receiving one standard therapy for SLE with any of the following: corticosteroids, hydroxychloroquine, or immunosuppressives (cyclophosphamide, azathioprine, mycophenolate, methotrexate, cyclosporine) AND there must be an absence of severe active lupus nephritis or severe active central nervous system lupus before Benlysta is authorized. B vs D coverage determination.

BEXAROTENE

Products Affected

- Bexarotene

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

BOTOX

Products Affected

- Botox

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Exclude when used for cosmetic purposes.
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial: 3 months. Continuation: 12 months
Other Criteria	<p>For chronic migraine, the patient has had failure or inadequate response following a minimum 3 month trial, contraindication per FDA label, intolerance, or not a candidate for at least TWO different prescription migraine prevention therapies from different classes of migraine prophylaxis medications: Antiepileptic drugs (ex: divalproex, sodium valproate, topiramate), Antidepressants (ex: amitriptyline, venlafaxine), Beta blockers (ex:metoprolol, propranolol, timolol, atenolol, nadolol)</p> <p>For overactive bladder, the patient has had failure or inadequate response two antimuscarinic medications for OAB (for example: darifenacin, flavoxate, oxybutynin, solifenacin, tolterodine, Toviaz).</p>

BRUKINSA

Products Affected

- Brukinsa

PA Criteria	Criteria Details
Indications	All FDA-approved indications not otherwise excluded from Part D
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Mantle Cell Lymphoma – approve for 3 years if the patient has tried at least one prior therapy.

BUPRENORPHINE

Products Affected

- Buprenorphine Hcl SUBL

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of opioid dependence.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Induction therapy: 1 month. Pregnancy/Hypersensitivity to naloxone: 12 months
Other Criteria	The use of buprenorphine for maintenance therapy should be limited to patients who have experienced a hypersensitivity reaction to naloxone or require buprenorphine during pregnancy.

CARBAGLU

Products Affected

- Carbaglu

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

CAYSTON

Products Affected

- Cayston

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of Cystic Fibrosis and documentation of Pseudomonas aeruginosa infection.
Age Restrictions	7 years of age and older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

CINRYZE

Products Affected

- Cinryze

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Patient must have a confirmed diagnosis of HAE.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	The patient must have a history of more than one severe event per month. B vs D coverage determination.

COLONY STIMULATING FACTORS

Products Affected

- Zarxio
- Neulasta
- Neulasta Onpro

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A

CORLANOR

Products Affected

- Corlanor TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Chronic stable angina, in combination with beta-blocker therapy
Exclusion Criteria	1.) Blood pressure less than 90/50 mmHg. 2.) Sick sinus syndrome, sinoatrial block, or 3rd degree AV block, unless a functioning demand pacemaker is present. 3.) Resting heart rate less than 60 bpm prior to treatment. 4.) Pacemaker dependence (heart rate maintained exclusively by the pacemaker).
Required Medical Information	Documentation of diagnosis, previous use of a beta-blocker, LVEF, sinus rhythm, resting HR, and blood pressure.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For use in chronic heart failure, the patient must have left ventricular ejection fraction less than or equal to 35%, normal sinus rhythm with resting heart rate greater than or equal to 70 beats per minute, and either be on maximally tolerated doses of beta-blockers or have a contraindication to beta-blocker use. For use in chronic angina, the patient must be using in combination with maximally tolerated doses of beta-blockers.

CYSTARAN

Products Affected

- Cystaran

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

DALFAMPRIDINE

Products Affected

- Dalfampridine Er

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Dalfampridine is considered medically necessary for patients with multiple sclerosis with medical documentation of impaired walking ability.

DEMSEER

Products Affected

- Demser

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For initial therapy, approve if the patient has tried a selective alpha blocker (e.g., doxazosin, terazosin or prazosin) AND the patient has tried phenoxybenzamine (brand or generic). For continuation of therapy, approve if the patient is currently receiving Demser or has received Demser in the past.

DERMATOLOGICAL RETINOIDS

Products Affected

- Avita
- Tretinoin CREA
- Tretinoin GEL
- Tretinoin Microsphere GEL 0.1%
- Tretinoin Microsphere Pump GEL 0.1%

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Coverage is not provided for cosmetic use.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

DERMATOLOGICAL WOUND CARE AGENTS

Products Affected

- Regranex

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

DIHYDROERGOTAMINE MESYLATE

Products Affected

- Dihydroergotamine Mesylate SOLN

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A

DRONABINOL

Products Affected

- Dronabinol

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	B vs D coverage determination

DUAVEE

Products Affected

- Duavee

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For the prevention of postmenopausal osteoporosis, trial, failure, or intolerance of raloxifene is required prior to the use of Duavee.

DUPIXENT

Products Affected

- Dupixent

PA Criteria	Criteria Details
Indications	All FDA approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Concurrent use with Xolair
Required Medical Information	Diagnosis, prescriber specialty, other medications tried and length of trials
Age Restrictions	asthma/AD-12 years of age and older. Chronic Rhinosinusitis-18 years of age and older
Prescriber Restrictions	Atopic Dermatitis-Prescribed by or in consultation with an allergist, immunologist or dermatologist, asthma-prescribed by or in consultation with an allergist, immunologist or pulmonologist
Coverage Duration	AD-Initial-16 weeks, Cont-1 year, asthma/Rhinosinusitis-initial-6 months, cont 1 year
Other Criteria	Atopic Dermatitis-Initial-meets both a and b: a.has used at least one medium, medium-high, high, and/or super-high-potency prescription topical corticosteroid OR has atopic dermatitis affecting ONLY the face, eyes/eyelids, skin folds, and/or genitalia and has tried tacrolimus ointment AND b.Inadequate efficacy was demonstrated with these previously tried topical prescription therapies, according to the prescribing physician.Continuation-Approve if the pt has responded to Dupixent therapy as determined by the prescribing physician. Asthma-Initial-approve if pt meets the following criteria (i, ii, and iii):i.Pt meets ONE of the following criteria (a or b):a)has a blood eosinophil level of greater than or equal to 150 cells per microliter within the previous 6 weeks or within 6 weeks prior to treatment with any anti-interleukin (IL) therapy or Xolair OR b)has oral corticosteroid-dependent asthma, per the prescriber AND ii.has received combination therapy with BOTH of the following (a and b): a)An inhaled corticosteroid (ICS) AND b)At least one additional asthma controller/maintenance medication (NOTE:An exception to the requirement for a trial of one additional asthma controller/maintenance medication can be made if the patient has already received anti-IL-5 therapy or Xolair used concomitantly with an ICS. Use of a combination inhaler containing both an ICS and a LABA would fulfil the requirement for both criteria a and b) AND iii.asthma is uncontrolled or was uncontrolled prior to starting any anti-IL therapy or Xolair as defined by ONE of the following (a, b, c, d or e): a)experienced two or more asthma

exacerbations requiring treatment with systemic corticosteroids in the previous year OR b)experienced one or more asthma exacerbation requiring hospitalization or an Emergency Department visit in the previous year OR c)has a forced expiratory volume in 1 second (FEV1) less than 80% predicted OR d)has an FEV1/forced vital capacity (FVC) less than 0.80 OR e)The patient's asthma worsens upon tapering of oral corticosteroid therapy. Continuation-Approve if meets the following criteria (i and ii): i.continues to receive therapy with one inhaled corticosteroid (ICS) or one ICS-containing combination inhaler AND ii.has responded to Dupixent therapy as determined by the prescribing physician. Chronic rhinosinusitis with Nasal Polyposis-Initial-pt is currently receiving therapy with an intranasal corticosteroid AND is experiencing significant rhinosinusitis symptoms such as nasal obstruction, rhinorrhea, or reduction/loss of smell according to the prescriber AND meets ONE of the following (a or b): a)has received treatment with a systemic corticosteroid within the previous 2 years or has a contraindication to systemic corticosteroid therapy OR b)has had prior surgery for nasal polyps. Continuation-approve if the pt continues to receive therapy with an intranasal corticosteroid AND pt has responded to Dupixent therapy as determined by the prescriber.

ENDOCRINE AND METABOLIC AGENTS

Products Affected

- Somavert

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient must have a diagnosis of acromegaly AND had inadequate response to surgery or radiation therapy or documentation these therapies are not appropriate for the patient.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

ENZYME REPLACEMENT/MODIFIERS

Products Affected

- Aldurazyme
- Elaprase
- Naglazyme

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

EPCLUSA

Products Affected

- Epclusa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation from the medical record of diagnosis including genotype, current medication regimen, HCV-RNA levels, history of previous HCV therapies and presence/absence of cirrhosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 weeks, based on indication and established treatment guidelines
Other Criteria	N/A

EPIDIOLEX

Products Affected

- Epidiolex

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of Lennox-Gastaut syndrome or Dravet syndrome
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

ERIVEDGE

Products Affected

- Erivedge

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of locally advanced or metastatic basal cell carcinoma that has recurred following surgery or who are not candidates for surgery, and who are not candidates for radiation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

ERLEADA

Products Affected

- Erleada

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Erleada is approved for use in combination with a gonadotropin-releasing hormone (GnRH) analog or in patients who have had a bilateral orchiectomy.

EVOMELA

Products Affected

- Evomela

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	B vs D coverage determination

EYLEA

Products Affected

- Eylea

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Administered by or under the supervision of an ophthalmologist
Coverage Duration	3 years
Other Criteria	N/A

FYCOMPA

Products Affected

- Fycompa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

GATTEX

Products Affected

- Gattex

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

HALAVEN

Products Affected

- Halaven

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For metastatic breast cancer, documentation of prior treatment with an anthracycline and a taxane. For unresectable or metastatic liposarcoma, documentation of prior treatment with an anthracycline-containing regimen.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For metastatic breast cancer, patients must have previously received at least two chemotherapeutic regimens for the treatment of metastatic disease. B vs D coverage determination.

HARVONI

Products Affected

- Harvoni

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation from the medical record of diagnosis including genotype, current medication regimen, HCV-RNA levels, history of previous HCV therapies and presence/absence of cirrhosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 to 24 weeks based on indication and established treatment guidelines
Other Criteria	N/A

HEMATOPOIETICS

Products Affected

- Aranesp Albumin Free INJ
100MCG/0.5ML, 100MCG/ML,
10MCG/0.4ML, 150MCG/0.3ML,
200MCG/0.4ML, 200MCG/ML,
25MCG/0.42ML, 25MCG/ML,
300MCG/0.6ML, 300MCG/ML,
40MCG/0.4ML, 40MCG/ML,
500MCG/ML, 60MCG/0.3ML,
60MCG/ML
- Retacrit

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For initial treatment of anemia, documentation of Hemoglobin less than 10, transferrin saturation greater than 20%, and ferritin levels greater than 100 obtained over the last 3 months. For patients who do not meet iron store requirements to create red blood cells, approval can be given if evidence shows the patient has started supplemental iron. For continuation of therapy, approvals granted if Hemoglobin does not exceed 11 g/dL for chronic kidney disease anemia, 13g/dL for the indication of reduction of allogeneic red cell transfusions in patients undergoing elective, noncardiac, nonvascular surgery, and 12g/dL for all other indications.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	B vs D coverage determination

HETLIOZ

Products Affected

- Hetlioz

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	A circadian rhythm longer than 24 hours has been confirmed by daily sleep logs and actigraphy for at least 14 days. Documentation that patient is totally blind and lacks light perception.
Age Restrictions	18 years of age and older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

HORMONAL AGENTS, GONADOTROPINS

Products Affected

- Chorionic Gonadotropin INJ

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

HORMONAL AGENTS, SOMATOSTATIN ANALOGS

Products Affected

- Octreotide Acetate
- Somatuline Depot

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

HRM - BENZTROPINE

Products Affected

- Benztropine Mesylate TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The physician has documented the indication for the continued use of the HRM (high risk med) with an explanation of the specific benefit established with the medication and how that benefit outweighs the potential risk, AND the physician will continue to monitor for side effects, AND the physician has documented that the patient has tried and failed two (2) safer formulary alternatives if two are available or provided clinical rationale why two safer formulary alternatives are not appropriate for the patient. If only one (1) safer formulary alternative is available, then only that particular medication would need to be documented as tried and failed or clinical rationale provided as to why that one safer formulary alternative is not appropriate for the patient. For patients concurrently taking multiple anticholinergic medications, the physician has assessed the risk.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior authorization required for age 65 or older.
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Safer alternatives depend on indication. For Parkinsonism, safer alternatives are: Carbidopa/Levodopa, Pramipexole, Ropinirole, Bromocriptine, Amantadine, and Selegiline. For extrapyramidal symptoms, a safer alternative is: Amantadine.

HRM - BUTALBITAL COMBINATIONS

Products Affected

- Butalbital/acetaminophen/caffeine CAPS
- Butalbital/acetaminophen/caffeine TABS 325MG; 50MG; 40MG
- Esgic CAPS
- Zebutal CAPS 325MG; 50MG; 40MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The physician has documented the indication for the continued use of the HRM (high risk med) with an explanation of the specific benefit established with the medication and how that benefit outweighs the potential risk, AND the physician will continue to monitor for side effects, AND the physician has documented that the patient has tried and failed two (2) safer formulary alternatives or provided clinical rationale why two safer formulary alternatives are not appropriate for the patient.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior authorization required for age 65 or older.
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Safer alternatives are: naproxen sodium and ibuprofen.

HRM - ESTROGENS

Products Affected

- Dotti
- Estradiol ORAL TABS 0.5MG, 1MG, 2MG
- Estradiol PTTW
- Estradiol PTWK
- Fyavolv
- Premarin TABS 0.3MG, 0.45MG, 0.625MG, 0.9MG, 1.25MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The physician has documented the indication for the continued use of the HRM (high risk med) with an explanation of the specific benefit established with the medication and how that benefit outweighs the potential risk, AND the physician will continue to monitor for side effects, AND the physician has documented that the patient has tried and failed two (2) safer formulary alternatives or provided clinical rationale why two safer formulary alternatives are not appropriate for the patient.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior authorization required for age 65 or older.
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Safer alternatives to the estrogen high risk medications depend on the indication. For Hot Flashes, safer alternatives are: SSRIs, venlafaxine, and gabapentin. For Vaginal Symptoms of menopause, safer alternatives are: Premarin Cream, Yuvafem and vaginal estradiol tablets. For Bone Density, safer alternatives are: bisphosphonates, raloxifene, and Prolia.

HRM - FIRST GENERATION ANTIHISTAMINES

Products Affected

- Hydroxyzine Hcl

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The physician has documented the indication for the continued use of the HRM (high risk med) with an explanation of the specific benefit established with the medication and how that benefit outweighs the potential risk, AND the physician will continue to monitor for side effects, AND the physician has documented that the patient has tried and failed two (2) safer formulary alternatives or provided clinical rationale why two safer formulary alternatives are not appropriate for the patient.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior authorization required for age 65 or older.
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Safer alternatives to hydroxyzine hydrochloride depend on the indication. Anxiety: buspirone, paroxetine, venlafaxine, escitalopram, sertraline, duloxetine. For all other indications, no safer alternatives are required.

HRM - MEGESTROL

Products Affected

- Megestrol Acetate SUSP 40MG/ML
- Megestrol Acetate TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The physician has documented the indication for the continued use of the HRM (high risk med) with an explanation of the specific benefit established with the medication and how that benefit outweighs the potential risk, AND the physician will continue to monitor for side effects.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior authorization required for age 65 or older.
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For the treatment of cachexia/loss of appetite associated with AIDS (megestrol oral suspension only), the physician has documented that the patient has tried and failed dronabinol or provided clinical rationale as to why that safer formulary alternative is not appropriate for the patient. For all other indications, trial of dronabinol is not required.

HRM - PERPHENAZINE/AMITRIPTYLINE

Products Affected

- Perphenazine/amitriptyline

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The physician has documented the indication for the continued use of the HRM (high risk med) with an explanation of the specific benefit established with the medication and how that benefit outweighs the potential risk, AND the physician will continue to monitor for side effects, AND the physician has documented that the patient has tried and failed two (2) safer formulary alternatives or provided clinical rationale why two safer formulary alternatives are not appropriate for the patient. For patients concurrently taking multiple anticholinergic medications, the physician has assessed the risk.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior authorization required for age 65 or older.
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Safer alternatives are: Nortriptyline, Protriptyline, Desipramine, Amoxapine, Citalopram, Venlafaxine, Fluoxetine, Paroxetine, Sertraline, Duloxetine, and Bupropion.

HRM - PLATELET MODIFYING AGENTS

Products Affected

- Dipyridamole TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The physician has documented the indication for the continued use of the HRM (high risk med) with an explanation of the specific benefit established with the medication and how that benefit outweighs the potential risk, AND the physician will continue to monitor for side effects, AND the physician has documented that the patient has tried and failed two (2) safer formulary alternatives or provided clinical rationale why two safer formulary alternatives are not appropriate for the patient.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior authorization required for age 65 or older.
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Safer alternatives are: Clopidogrel, Warfarin, Jantoven, and aspirin/dipyridamole .

HRM - PROMETHAZINE

Products Affected

- Promethazine Hcl SYRP
- Promethazine Hcl TABS 12.5MG
- Promethazine Hcl Plain
- Promethazine Hydrochloride TABS
25MG, 50MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The physician has documented the indication for the continued use of the HRM (high risk med) with an explanation of the specific benefit established with the medication and how that benefit outweighs the potential risk, AND the physician will continue to monitor for side effects, AND the physician has documented that the patient has tried and failed two safer formulary alternatives if available or provided clinical rationale why the safer formulary alternative is not appropriate for the patient. If only one (1) safer formulary alternative is available, then only that particular medication would need to be documented as tried and failed or clinical rationale provided as to why that one safer formulary alternative is not appropriate for the patient. For patients concurrently taking other anticholinergic medications, the physician has assessed the risk.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior authorization required for age 65 or older.
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For postoperative nausea and vomiting, the safer alternative is ondansetron. For perennial and seasonal allergic rhinitis, safer alternatives are: levocetirizine and desloratadine. For any other indications, trial of a formulary alternative is not required.

HRM - SKELETAL MUSCLE RELAXANTS

Products Affected

- Cyclobenzaprine Hydrochloride
TABS 10MG, 5MG
- Methocarbamol TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The physician has documented the indication for the continued use of the HRM (high risk med) with an explanation of the specific benefit established with the medication and how that benefit outweighs the potential risk, AND the physician will continue to monitor for side effects. For patients concurrently taking multiple anticholinergic medications, the physician has assessed the risk.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior authorization required for age 65 or older.
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

HRM - TRICYCLIC ANTIDEPRESSANTS

Products Affected

- Amitriptyline Hcl TABS 100MG, 150MG, 25MG, 75MG
- Amitriptyline Hydrochloride TABS 10MG, 50MG
- Clomipramine Hcl CAPS
- Doxepin Hcl CAPS 100MG, 10MG, 150MG, 50MG, 75MG
- Doxepin Hcl CONC
- Doxepin Hydrochloride CAPS 25MG
- Imipramine Hcl TABS 25MG, 50MG
- Imipramine Hydrochloride TABS 10MG
- Trimipramine Maleate CAPS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The physician has documented the indication for the continued use of the HRM (high risk med) with an explanation of the specific benefit established with the medication and how that benefit outweighs the potential risk, AND the physician will continue to monitor for side effects AND the physician has documented that the patient has tried and failed two (2) safer formulary alternatives or provided clinical rationale why two safer formulary alternatives are not appropriate for the patient. For patients concurrently taking multiple anticholinergic medications, the physician has assessed the risk.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior authorization required for age 65 or older.
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Safer alternatives depend on indication. For Depression, safer alternatives are: Nortriptyline, Protriptyline, Desipramine, Amoxapine, Citalopram, Escitalopram, Fluvoxamine, Venlafaxine, Fluoxetine, Paroxetine, Sertraline, Duloxetine, and Bupropion. For headache prophylaxis, safer alternatives are: Propranolol, Topiramate, and Divalproex sodium. For headache treatment, safer alternatives are: Sumatriptan, Naratriptan, Rizatriptan and Dihydroergotamine. For Pain/Neuropathy, safer alternatives are: Duloxetine, Lyrica, and Gabapentin. If using requested medication for a medically-accepted indication not listed above, then no trial of alternatives is required.

HYDROXYPROGESTERONE

Products Affected

- Hydroxyprogesterone Caproate INJ
1.25GM/5ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	B vs D coverage determination

ICATIBANT

Products Affected

- Icatibant Acetate

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Patient must have a confirmed diagnosis of HAE.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Icatibant is considered medically necessary for the treatment of acute attacks of hereditary angioedema (HAE) in patients who have tried and failed Ruconest.

IDIOPATHIC PULMONARY FIBROSIS

Products Affected

- Esbriet
- Ofev

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	For the diagnosis of IPF, other known causes of interstitial lung disease e.g., domestic and occupational environmental exposures, connective tissue disease, and drug toxicity.
Required Medical Information	Diagnosis of IPF confirmed by 1.) in patients without surgical lung biopsy: Usual interstitial pneumonia (UIP) pattern on high resolution computed tomography (HRCT) is indicative of IPF or 2.) in patients with surgical lung biopsy: The biopsy pattern is diagnostic of IPF or the combination of HRCT and biopsy pattern is indicative of IPF. For use of Ofev in systemic sclerosis-associated interstitial lung disease (SSc-ILD) no further documentation is required.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Esbriet and Ofev will each be used as monotherapy.

IMMUNE STIMULANTS

Products Affected

- Adagen

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

IMMUNE SUPPRESSANTS

Products Affected

- Enbrel
- Enbrel Mini
- Enbrel Sureclick
- Humira
- Humira Pediatric Crohns Disease Starter Pack
- Humira Pen
- Humira Pen-cd/uc/hs Starter
- Humira Pen-ps/uv Starter
- Skyrizi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis and past medication history.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	<p>1.) Use of Humira or Enbrel is considered medically necessary for the treatment of Rheumatoid Arthritis in patients that have tried and failed methotrexate OR at least one alternative disease modifying antirheumatic drugs (DMARDs). A previous trial of a biologic also counts as a trial.</p> <p>2.) Use of Humira or Enbrel is considered medically necessary for the treatment of Juvenile Rheumatoid Arthritis in patients that have tried and failed at least one other agent (e.g. MTX, sulfasalazine, leflunomide, or DMARD) or will be starting on a adalimumab concurrently with MTX, sulfasalazine, or leflunomide. Approve without trying another agent if patient has absolute contraindication to MTX, sulfasalazine, or leflunomide or if patient has aggressive disease. A previous trial of a biologic also counts as a trial.</p> <p>3.) Use of Humira or Enbrel is considered medically necessary for the treatment of Ankylosing Spondylitis in patients that have tried and failed at least 1 non-steroidal anti-inflammatory drug (NSAID), corticosteroid, OR sulfasalazine. A previous trial of a biologic also counts as a trial.</p> <p>4.) Use of Humira, Enbrel, or Sykrizi is considered medically necessary for the treatment of Plaque Psoriasis in patients that have: a.) moderate to severe chronic disease, b.) minimum body surface area (BSA) involvement of greater than or equal to 5% OR involvement of the palms, soles, head, neck or genitalia, AND c.) tried and failed at least 1 topical agent (topical steroid, calcipotriene, or tazarotene) OR one systemic agent (e.g. MTX, cyclosporine, etc.) A previous trial of a biologic also counts as a trial.</p>

-
- 5.) Use of Humira or Enbrel is considered medically necessary for the treatment of Psoriatic Arthritis in patients with active disease.
- 6.) Use of Humira is considered medically necessary for the treatment of moderate to severe Crohn's Disease in patients that have tried and failed at least 1 of the following: immunomodulators, corticosteroids, or aminosalicylates. A previous trial of a biologic also counts as a trial.
- 7.) Use of Humira is considered medically necessary for the treatment of moderately to severely active ulcerative colitis in patients who have had inadequate response to at least 1 of the following: corticosteroids, sulfasalazine, mesalamine, azathioprine, 6-mercaptopurine, cyclosporine, tacrolimus. A previous trial of a biologic also counts as a trial.
- 8.) Use of Humira is considered medically necessary for the treatment of hidradenitis suppurativa and uveitis.
-

IMMUNE SUPPRESSANTS - RINVOQ

Products Affected

- Rinvoq

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis and past medication history.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	1.) Rinvoq is considered medically necessary for the treatment of Rheumatoid Arthritis in patients that have tried and failed methotrexate OR at least one alternative disease modifying antirheumatic drugs (DMARDs). A previous trial of a biologic also counts as a trial.

IMMUNE SUPPRESSANTS - STELARA

Products Affected

- Stelara INJ 45MG/0.5ML, 90MG/ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis and past medication history.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	<p>1.) Stelara is considered medically necessary for the treatment of Plaque Psoriasis in patients that have: a.) moderate to severe chronic disease, b.) minimum body surface area (BSA) involvement of greater than or equal to 5% OR involvement of the palms, soles, head, neck or genitalia, AND c.) tried and failed at least 1 topical agent (topical steroid, calcipotriene, or tazarotene) OR one systemic agent (e.g. MTX, cyclosporine, etc.). A previous trial of a biologic also counts as a trial.</p> <p>2.) Stelara is considered medically necessary for the treatment of Psoriatic Arthritis in patients with active disease.</p> <p>3.) Stelara is considered medically necessary for the treatment of moderate to severe Crohn's Disease in patients that have tried and failed at least 1 of the following: immunomodulators, corticosteroids, or aminosalicylates. A previous trial of a biologic also counts as a trial.</p> <p>4.) Stelara is considered medically necessary for the treatment of moderately to severely active ulcerative colitis in patients who have had inadequate response to at least 1 of the following: corticosteroids, sulfasalazine, mesalamine, azathioprine, 6-mercaptopurine, cyclosporine, tacrolimus. A previous trial of a biologic also counts as a trial.</p>

IMMUNE SUPPRESSANTS - TRANSPLANT RELATED

Products Affected

- Astagraf XL
- Azathioprine INJ
- Azathioprine TABS
- Cyclosporine CAPS
- Cyclosporine INJ
- Cyclosporine Modified
- Gengraf CAPS 100MG, 25MG
- Gengraf SOLN
- Mycophenolate Mofetil
- Mycophenolic Acid Dr
- Prograf PACK
- Rapamune SOLN
- Sandimmune SOLN
- Sirolimus SOLN
- Sirolimus TABS
- Tacrolimus CAPS
- Zortress

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	B vs D coverage determination

IMMUNE SUPPRESSANTS - XELJANZ

Products Affected

- Xeljanz
- Xeljanz Xr

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis and past medication history.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	<p>1.) Xeljanz is considered medically necessary for the treatment of Rheumatoid Arthritis in patients that have tried and failed methotrexate OR at least one alternative disease modifying antirheumatic drugs (DMARDs). A previous trial of a biologic also counts as a trial. For Rheumatoid Arthritis, dosing 10mg twice a day will not be approved in patients with at least one cardiovascular risk factor.</p> <p>2.) Xeljanz is considered medically necessary for the treatment of Psoriatic Arthritis in patients with active disease. For Psoriatic Arthritis, dosing 10mg twice a day will not be approved in patients with at least one cardiovascular risk factor.</p> <p>3.) Xeljanz is considered medically necessary for the treatment of moderately to severely active ulcerative colitis in patients who have had inadequate response to at least 1 of the following: corticosteroids, sulfasalazine, mesalamine, azathioprine, 6-mercaptopurine, cyclosporine, tacrolimus. A previous trial of a biologic also counts as a trial.</p>

IMMUNOMODULATORS

Products Affected

- Avonex
- Avonex Pen
- Betaseron
- Copaxone INJ 20MG/ML, 40MG/ML
- Gilenya CAPS 0.5MG
- Tecfidera
- Tecfidera Starter Pack

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Multiple Sclerosis, Clinically Isolated Syndrome (Avonex and Betaseron)
Exclusion Criteria	N/A
Required Medical Information	Documentation of a relapsing form of multiple sclerosis (MS) (e.g., relapsing-remitting MS, progressive-relapsing MS, or secondary progressive MS with relapses), or diagnosis of Clinically Isolated Syndrome.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Requests for Clinically Isolated Syndrome will be approved for Avonex and Betaseron only.

INFLIXIMAB

Products Affected

- Renflexis

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis and past medication history.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	<p>1.) Use of Renflexis is considered medically necessary for the treatment of Rheumatoid Arthritis in patients that have tried and failed methotrexate OR at least one alternative disease modifying antirheumatic drugs (DMARDs).</p> <p>2.) Use of Renflexis is considered medically necessary for the treatment of Juvenile Rheumatoid Arthritis in patients that have tried and failed at least one other agent (e.g. MTX, sulfasalazine, leflunomide, or DMARD). Approve without trying another agent if patient has absolute contraindication to MTX, sulfasalazine, or leflunomide or if patient has aggressive disease.</p> <p>3.) Use of Renflexis is considered medically necessary for the treatment of Ankylosing Spondylitis in patients that have tried and failed at least 1 non-steroidal anti-inflammatory drug (NSAID), corticosteroid, OR sulfasalazine.</p> <p>4.) Use of Renflexis is considered medically necessary for the treatment of Plaque Psoriasis in patients that have: a.) moderate to severe chronic disease, b.) minimum body surface area (BSA) involvement of greater than or equal to 5% OR involvement of the palms, soles, head, neck or genitalia, AND c.) tried and failed at least 1 topical agent (topical steroid, calcipotriene, or tazarotene) OR one systemic agent (e.g. MTX, cyclosporine, etc.)</p> <p>5.) Use of Renflexis is considered medically necessary for the treatment of Psoriatic Arthritis in patients with active disease.</p> <p>6.) Use of Renflexis is considered medically necessary for the treatment of moderate to severe Crohn's Disease in patients that have tried and</p>

failed at least 2 of the following: immunomodulators, corticosteroids, or aminosalicylates

7.) Use of Renflexis is considered medically necessary for the treatment of moderately to severely active ulcerative colitis in patients who have had inadequate response to at least 2 of the following: corticosteroids, sulfasalazine, mesalamine, azathioprine, 6-mercaptopurine.

8.) Use of Renflexis is considered medically necessary for the treatment of fistulizing Crohn's disease.

B vs D coverage determination required for Renflexis.

INSULIN-LIKE GROWTH FACTOR

Products Affected

- Increlex

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis. Documentation of lab data reflecting height standard deviation score, basal IGF-1 score, and growth hormone level.
Age Restrictions	Patients 2 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Height standard deviation score must be less than or equal to -3.0 AND the basal IGF-1 score must be below the lower limits of normal for the reporting lab AND the patient must have a normal or elevated growth hormone level (excluding patients with growth hormone gene deletion) AND epiphyses must be confirmed as open in patients greater than or equal to 10 years of age.

ISTODAX

Products Affected

- Istodax (overfill)
- Romidepsin

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis and past medication history.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Use of Istodax is considered medically necessary for the treatment of cutaneous T-cell lymphoma in patients that have tried and failed at least 1 prior therapy. B vs D coverage determination.

JADENU

Products Affected

- Jadenu

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	<p>Transfusion-related chronic iron overload, initial therapy - approve if the patient is receiving blood transfusions at regular intervals for various conditions (eg, thalassemia syndromes, myelodysplastic syndrome, chronic anemia, sickle cell disease) AND prior to starting therapy, the serum ferritin level is greater than 1,000 mcg/L.</p> <p>Non-transfusion-dependent thalassemia syndromes chronic iron overload, initial therapy - approve if prior to starting therapy the serum ferritin level is greater than 300 mcg/L.</p> <p>Continuation therapy - approve if the patient is benefiting from therapy as confirmed by the prescribing physician.</p>

KALYDECO

Products Affected

- Kalydeco

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Patients with cystic fibrosis (CF) who are homozygous for the F508del mutation in the CFTR gene. Kalydeco will not be used in combination with Orkambi or Symdeko
Required Medical Information	CF mutation test documenting patient has one mutation in the CFTR gene that is responsive to ivacaftor based on clinical and/or in vitro assay data.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

KORLYM

Products Affected

- Korlym

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years of age and older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

KUVAN

Products Affected

- Kuvan

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis, Phe concentration
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial Approval: 2 months. Continuation of therapy: 12 months.
Other Criteria	For continuation of therapy: the patient must have responded to a therapeutic trial of Kuvan. Response is defined as a 20% or greater reduction in blood phenylalanine level from baseline.

LIDOCAINE PATCH

Products Affected

- Lidocaine PTCH

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For the FDA-labeled indication of post-herpetic neuralgia, no additional criteria are required to be met. For diabetic neuropathic pain: the patient must have previous use and inadequate response or intolerance to any ONE medication that is FDA-labeled for diabetic peripheral neuropathy, including duloxetine and Lyrica. For cancer related neuropathic pain (including treatment-related neuropathy), no additional criteria are required to be met.

LUMIZYME

Products Affected

- Lumizyme

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	B vs D coverage determination

MAVYRET

Products Affected

- Mavyret

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation from the medical record of diagnosis including genotype, current medication regimen, HCV-RNA levels, history of previous HCV therapies and presence/absence of cirrhosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	8-16 weeks, based on indication and established treatment guidelines
Other Criteria	N/A

METABOLIC BONE DISEASE AGENTS

Products Affected

- Forteo INJ 600MCG/2.4ML
- Tymlos

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Cumulative use of parathyroid hormone analogs for more than 2 years during a patient's lifetime
Required Medical Information	Diagnosis of osteoporosis based on DEXA (T-score less than or equal to -2.5) or based on presence of documented fragility fracture.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	2 years from initiation of therapy
Other Criteria	Member has tried and/or failed a bisphosphonate or SERM OR the member has documented intolerance, contraindication, or hypersensitivity to other osteoporosis therapies. For patients with a T-score less than or equal to -3.5, failure of bisphosphonates or SERMs are not required.

MOLECULAR TARGET INHIBITORS

Products Affected

- Abiraterone Acetate
- Afinitor
- Afinitor Disperz
- Alecensa
- Alunbrig
- Balversa
- Bosulif
- Braftovi CAPS 75MG
- Cabometyx
- Calquence
- Caprelsa
- Cometriq
- Copiktra
- Cotellic
- Daurismo
- Erlotinib Hydrochloride
- Everolimus (antineoplastic)
- Farydak
- Gilotrif
- Ibrance
- Iclusig
- Idhifa
- Imatinib Mesylate
- Imbruvica
- Inlyta
- Inrebic
- Iressa
- Jakafi
- Kisqali
- Kisqali Femara 200 Dose
- Kisqali Femara 400 Dose
- Kisqali Femara 600 Dose
- Lenvima 10 Mg Daily Dose
- Lenvima 12mg Daily Dose
- Lenvima 14 Mg Daily Dose
- Lenvima 18 Mg Daily Dose
- Lenvima 20 Mg Daily Dose
- Lenvima 24 Mg Daily Dose
- Lenvima 4 Mg Daily Dose
- Lenvima 8 Mg Daily Dose
- Lonsurf
- Lorbreana

- Lynparza TABS
- Mekinist
- Mektovi
- Nerlynx
- Nexavar
- Ninlaro
- Piqray 200mg Daily Dose
- Piqray 250mg Daily Dose
- Piqray 300mg Daily Dose
- Pomalyst
- Rubraca
- Rydapt
- Sprycel
- Stivarga
- Sutent
- Synribo
- Tafinlar
- Tagrisso
- Talzenna
- Tassigna
- Tibsovo
- Tykerb
- Venclexta
- Venclexta Starting Pack
- Verzenio
- Vizimpro
- Votrient
- Xalkori
- Xospata
- Xpovio 100 Mg Once Weekly
- Xpovio 60 Mg Once Weekly
- Xpovio 80 Mg Once Weekly
- Xpovio 80 Mg Twice Weekly
- Xtandi
- Zejula
- Zelboraf
- Zydelig
- Zykadia
- Zytiga TABS 500MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

MONOCLONAL ANTIBODIES

Products Affected

- Xolair

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For the diagnosis of asthma: Laboratory data reflecting IgE levels greater than 30 IU/mL, medical history documenting previous trial and inadequate response to inhaled corticosteroids and a second controller such as a long-acting beta-agonist or a leukotriene receptor antagonist. For the diagnosis of chronic idiopathic urticaria (CIU): Documentation that the patient has remained symptomatic despite at least 4 weeks of a second generation H1 antihistamine (such as but not limited to levoceterizine or desloratadine) therapy at twice the recommended dosing.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Xolair will not be used concomitantly with Cinqair, Dupixent, Fasenna or Nucala

NATPARA

Products Affected

- Natpara

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Chronic hypoparathyroidism, initial therapy - approve if before starting Natpara, serum calcium concentration is greater than 7.5 mg/dL and 25-hydroxyvitamin D stores are sufficient per the prescribing physician. Chronic hypoparathyroidism, continuing therapy - approve if during Natpara therapy, the patient's 25-hydroxyvitamin D stores are sufficient per the prescribing physician AND the patient is responding to Natapara therapy, as determined by the prescriber.

NAYZILAM

Products Affected

- Nayzilam

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Use will be for treatment of intermittent, stereotypic episodes of frequent seizure activity (for example, seizure clusters, acute repetitive seizures) that are distinct from the individual's usual seizure pattern.
Age Restrictions	12 years of age and older
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Individual is currently receiving maintenance antiepileptic medication.

NMDA RECEPTOR ANTAGONIST

Products Affected

- Memantine Hcl Titration Pak
- Memantine Hydrochloride SOLN
- Memantine Hydrochloride TABS
- Memantine Hydrochloride Er
- Namzaric

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Automatic approval if member is greater than 26 years of age. Prior Authorization is required for age 26 or younger.
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

NON-AMPHETAMINE CENTRAL NERVOUS SYSTEM AGENTS

Products Affected

- Armodafinil

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis and sleep study for the diagnosis of sleep apnea or narcolepsy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

NORTHERA

Products Affected

- Northera

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Northera will be used for patients with neurogenic orthostatic hypotension associated with one of the following diagnoses: 1.) Primary autonomic failure due to Parkinson's disease, multiple system atrophy, or pure autonomic failure, OR 2.) Dopamine beta hydroxylase deficiency, OR 3.) Non-diabetic autonomic neuropathy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	The patient must have failure, contraindication or intolerance to fludrocortisone acetate or midodrine.

NUEDEXTA

Products Affected

- Nuedexta

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

NULOJIX

Products Affected

- Nulojix

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of Epstein-Barr virus serology and current medication regimen.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Documentation of use in combination with basiliximab induction, mycophenolate mofetil, and corticosteroids. B vs D coverage determination.

NUPLAZID

Products Affected

- Nuplazid

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For the treatment of Parkinson's disease psychosis: The patient has experienced hallucinations or delusions associated with Parkinson's disease psychosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

Ocaliva

Products Affected

- Ocaliva

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prescriber specialty, lab values, prior medications used for diagnosis and length of trials
Age Restrictions	18 years and older (initial and continuation therapy)
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant physician (initial and continuation therapy)
Coverage Duration	6 months initial, 3 years cont.
Other Criteria	Initial treatment of PBC-Patient must meet both 1 and 2-1. Patient has a diagnosis of PBC as defined by TWO of the following:a)Alkaline phosphatase (ALP) elevated above the upper limit of normal as defined by normal laboratory reference values b)Positive anti-mitochondrial antibodies (AMAs) c)Histologic evidence of primary biliary cholangitis (PBC) from a liver biopsy 2. Patient meets ONE of the following: a) Patient has been receiving ursodiol therapy for greater than or equal to 1 year and has had an inadequate response. b) Patient is unable to tolerate ursodiol therapy. Cont tx - approve if the patient has responded to Ocaliva therapy as determined by the prescribing physician (e.g., improved biochemical markers of PBC (e.g., alkaline phosphatase [ALP], bilirubin, gamma-glutamyl transpeptidase [GGT], aspartate aminotransferase [AST], alanine aminotransferase [ALT] levels)).

OCREVUS

Products Affected

- Ocrevus

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by, or in consultation with, a physician who specializes in the treatment of MS and/or a neurologist
Coverage Duration	1 year
Other Criteria	N/A

ODOMZO

Products Affected

- Odomzo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For the treatment of Locally Advanced Basal Cell Carcinoma: the cancer has recurred following surgery or radiation therapy OR the patient is not a candidate for surgery or radiation therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

ORENCIA

Products Affected

- Orenzia INJ 125MG/ML, 50MG/0.4ML, 87.5MG/0.7ML
- Orenzia Clickject

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis and past medication history.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	<p>1.) Orenzia SC is considered medically necessary for the treatment of Rheumatoid Arthritis in patients that have a.) tried and failed methotrexate OR an alternative disease modifying antirheumatic drugs (DMARDs) AND b.) tried and failed Enbrel or Humira.</p> <p>2.) Use of Orenzia SC is considered medically necessary for the treatment of Juvenile Rheumatoid Arthritis in patients that have a.) tried and failed at least 1 DMARD AND b.) tried and failed Enbrel or Humira.</p> <p>3.) Use of Orenzia SC is considered medically necessary for the treatment of: Psoriatic Arthritis in patients with active disease that have tried and failed Enbrel or Humira.</p>

ORKAMBI

Products Affected

- Orkambi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Combination use with Symdeko
Required Medical Information	CF mutation test documenting the patient is homozygous for the F508del mutation in the CFTR gene.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

PADCEV

Products Affected

- Padcev

PA Criteria	Criteria Details
Indications	All FDA-approved indications not otherwise excluded from Part D.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies
Age Restrictions	18 years and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	1 year
Other Criteria	Urothelial carcinoma-approve if the patient has locally advanced or metastatic disease, has previously received a programmed death receptor-1 or programmed death-ligand 1 inhibitor and has previously received platinum containing chemotherapy. B vs D coverage determination

PHOSPHODIESTERASE TYPE 4 (PDE4) INHIBITORS

Products Affected

- Daliresp

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For initial therapy: trial and failure, contraindication, or intolerance to two formulary inhaled COPD agents is required. For continuation of therapy, approve if patient has already been started on Daliresp.

PITUITARY HORMONES

Products Affected

- Genotropin
- Genotropin Miniquick
- Leuprolide Acetate INJ
- Lupron Depot (1-month)
- Lupron Depot (3-month)
- Lupron Depot (4-month)
- Lupron Depot (6-month)
- Lupron Depot-ped (1-month)
- Lupron Depot-ped (3-month)
- Nubeqa
- Synarel
- Trelstar Mixject
- Triptodur

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

PROMACTA

Products Affected

- Promacta

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Use of Promacta for the treatment of thrombocytopenia is considered medically necessary in: a.) patients with chronic hepatitis C, or b.) patients with chronic immune (idiopathic) thrombocytopenic purpura (ITP) that have failed corticosteroid OR intravenous immune globulin (IVIG) therapy OR have had an insufficient response to a splenectomy, or c.) severe aplastic anemia with documentation of inadequate response to previous immunosuppressive therapy (e.g. Atgam, Thymoglobulin, cyclosporine) or being used in combination with standard immunosuppressive therapy.

PURIXAN

Products Affected

- Purixan

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Documentation of trial, contraindication, or failure to mercaptopurine tablets.

QUNINIE SULFATE

Products Affected

- Quinine Sulfate CAPS 324MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Babesiosis, uncomplicated Plasmodium vivax malaria.
Exclusion Criteria	Excluded if used for treatment or prevention of nocturnal leg cramps.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 month
Other Criteria	N/A

RELISTOR

Products Affected

- Relistor INJ

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis and past medication history.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	Use of Relistor is considered medically necessary for the treatment of opioid-induced constipation in patients with advanced illnesses who are receiving palliative care AND have tried and failed laxative therapy with lactulose or polyethylene glycol. Relistor is also considered medically necessary for the treatment of opioid-induced constipation in adult patients with chronic non-cancer pain who have tried and failed laxative therapy with lactulose or polyethylene glycol AND Amitiza.

REPATHA

Products Affected

- Repatha
- Repatha Pushtronex System
- Repatha Sureclick

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of one of the following: Clinical atherosclerotic cardiovascular disease (ASCVD) OR Primary Hyperlipidemia (including Heterozygous familial hypercholesterolemia (HeFH)) OR Homozygous Familial Hypercholesterolemia (HoFH) (confirmed by either: Genetic testing or History of untreated low-density lipoprotein cholesterol (LDL-C) greater than 500mg/dl or treated LDL greater than 300mg/dl with either: presence of xanthomas before the age of 10 years or evidence of heterozygous familial hypercholesterolemia in both parents.)
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial:6 mo. Cont. of therapy: 12 months
Other Criteria	For ASCVD or Primary Hyperlipidemia (including HeFH): Patient is on high-intensity statin therapy or maximally tolerated statin therapy, is not at LDL-C level goal (LDL-C greater than 70 mg/dL) and will be continued on high intensity or maximally tolerated statin therapy while on the PCSK9-inhibitor. Patients intolerant to statins as demonstrated by experiencing statin-associated rhabdomyolysis to one statin OR has tried 2 statins (any combination of high or moderate intensity statins) and has experienced skeletal–muscle related symptoms on both agents, no concurrent statin use required. For HoFH: Patient is on high-intensity or maximally tolerated lipid lowering therapy (such as statins and/or ezetimibe), is not at LDL-C level goal (LDL-C greater than 70 mg/dL), and will be continued on high intensity or maximally tolerated lipid lowering therapy while on Repatha. For patients with HoFH who are intolerant to statins, no concurrent statin use required. Continuation of therapy requires documented evidence of clinical benefit response.

REVLIMID

Products Affected

- Revlimid

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

ROZLYTREK

Products Affected

- Rozlytrek

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	1. Documentation of metastatic non-small cell lung cancer (NSCLC) whose tumors are ROS1-positive. OR 2. Documentation of a solid tumors that has a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation. Patient is metastatic or where surgical resection is likely to result in severe morbidity. Patient has no satisfactory alternative treatments or has progressed following treatment.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A

RUCONEST

Products Affected

- Ruconest

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Patient must have a confirmed diagnosis of HAE.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

SAMSCA

Products Affected

- Samsca

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Maximum of 30 days for each course of treatment (initial or retreatment)
Other Criteria	Samsca is considered medically necessary for the treatment of patients with significant hypovolemic and euvolemic hyponatremia (serum sodium less than 125 mEq/L) or symptomatic hyponatremia.

SIGNIFOR

Products Affected

- Signifor

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

SIRTURO

Products Affected

- Sirturo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation from the medical record required indicating the patient has multi-drug resistant tuberculosis resistant to isoniazid and rifampin.
Age Restrictions	18 years of age and older
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	Use of Sirturo for the treatment of multi-drug resistant tuberculosis is considered medically necessary in patients with multi-drug resistant tuberculosis in combination with at least 3 other agents.

SODIUM PHENYL BUTYRATE

Products Affected

- Sodium Phenylbutyrate POWD
3GM/TSP
- Sodium Phenylbutyrate TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

SYLATRON

Products Affected

- Sylatron

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

SYMPAZAN

Products Affected

- Sympazan

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Failure, contraindication, or intolerance to clobazam tablets and oral solution before Sympazan is authorized.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

SYNAGIS

Products Affected

- Synagis INJ 100MG/ML,
50MG/0.5ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	24 months or younger
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A

TARGRETIN (TOPICAL)

Products Affected

- Targretin GEL

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

TETRABENAZINE

Products Affected

- Tetrabenazine

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis of chorea associated with Huntington's Disease. CYP 2D6 genotype must be provided for doses greater than 50mg/day.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

THALIDOMIDE (THALOMID)

Products Affected

- Thalomid

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

THIOTEPA

Products Affected

- Thiotepa INJ 15MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	B vs D coverage determination

TRANSMUCOSAL FENTANYL CITRATE

Products Affected

- Fentanyl Citrate Oral Transmucosal

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation from the medical record of diagnosis.
Age Restrictions	16 years of age and older for fentanyl citrate (lozenge/troche)
Prescriber Restrictions	Enrollment in the Transmucosal Immediate-Release Fentanyl (TIRF) REMS Access program.
Coverage Duration	12 months
Other Criteria	Transmucosal fentanyl products will only be covered with documentation of breakthrough cancer pain. The patient must be currently receiving and be tolerant to opioid therapy for persistent cancer pain. The patient must be enrolled in the TIRF REMS Access program.

TYSABRI

Products Affected

- Tysabri

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	<p>Treatment of relapsing forms of multiple sclerosis (MS) when EITHER of the following criteria is met: 1.) history of beneficial clinical response to Tysabri (natalizumab) for MS or 2.) failure, contraindication or intolerance to one formulary alternative (eg. Avonex, Betaseron, Copaxone, Gilenya, or Tecfidera).</p> <p>Treatment of moderately to severely active Crohn's disease (CD) when EITHER of the following criteria is met: 1.) history of beneficial clinical response to Tysabri (natalizumab) for CD or 2.) failure or intolerance to Humira.</p>

UPTRAVI

Products Affected

- Uptravi

PA Criteria	Criteria Details
Indications	All FDA approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of pulmonary arterial hypertension.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Patient new to Uptravi therapy must meet a) OR b): a) tried TWO or is currently taking TWO oral therapies for PAH (either alone or in combination) each for 30 days, unless patient has experienced treatment failure, intolerance, or oral therapy is contraindicated: PDE5 inhibitor (eg, sildenafil, Revatio), endothelin receptor antagonist (ERA) [eg, Tracleer, Letairis or Opsumit], or Adempas, OR b) receiving or has received in the past one prostacyclin therapy for PAH (eg, Orenitram, Ventavis, or epoprostenol injection).

VALCHLOR GEL

Products Affected

- Valchlor

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis and past medical history.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Valchlor Topical Gel is considered medically necessary for the treatment of patients with Stage 1A and 1B mycosis fungoides-type cutaneous T-cell lymphoma who have received prior skin-directed therapy.

VASODILATORS

Products Affected

- Adempas
- Ambrisentan
- Letairis
- Opsumit
- Sildenafil Citrate TABS 20MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Secondary Reynaud's phenomenon (sildenafil), Rebound pulmonary hypertension caused by nitric oxide withdrawal (sildenafil), Persistent pulmonary hypertension of the newborn (sildenafil)
Exclusion Criteria	N/A
Required Medical Information	Documentation of pulmonary arterial hypertension. Adempas will also be approved for chronic thromboembolic pulmonary hypertension.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Sildenafil will also be considered medically necessary for Secondary Reynaud's phenomenon, rebound pulmonary hypertension caused by nitric oxide withdrawal, or persistent pulmonary hypertension of the newborn.

VENTAVIS

Products Affected

- Ventavis

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	B vs D coverage determination

VIBERZI

Products Affected

- Viberzi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of irritable bowel syndrome with diarrhea.
Age Restrictions	18 years of age and older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	The patient must have a history of failure, contraindication or intolerance to one antidiarrheal drug.

VIGABATRIN

Products Affected

- Vigabatrin
- Vigadrone

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation from the medical record of diagnosis and past medication history.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Vigabatrin is considered medically necessary in patients that have failed to receive a clinically appropriate response from optimal doses and administration of at least two of the following: phenytoin, divalproex, lamotrigine, and levetiracetam. For the indication of Infantile Spasms, failure of another drug(s) is not required.

VITRAKVI

Products Affected

- Vitrakvi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of a solid tumors that has a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation. Patient is metastatic or where surgical resection is likely to result in severe morbidity. Patient has no satisfactory alternative treatments or has progressed following treatment.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

VIVITROL

Products Affected

- Vivitrol

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Documentation of a solid tumors that has a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation. Patient is metastatic or where surgical resection is likely to result in severe morbidity. Patient has no satisfactory alternative treatments or has progressed following treatment.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

VOSEVI

Products Affected

- Vosevi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Treatment naive patients (exception: treatment-naïve patients with genotype 3 with compensated cirrhosis and a Y93H mutation will be approved)
Required Medical Information	Documentation from the medical record of diagnosis including genotype, current medication regimen, HCV-RNA levels, history of previous HCV therapies and presence/absence of cirrhosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 weeks, based on indication and established treatment guidelines
Other Criteria	N/A

XATMEP

Products Affected

- Xatmep

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

XGEVA

Products Affected

- Xgeva

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For initial therapy in the treatment of hypercalcemia of malignancy: trial and failure, contraindication, or intolerance to one intravenous bisphosphonate (e.g. pamidronate, zoledronic acid) is required. For continuation of therapy in the treatment of hypercalcemia of malignancy, approve if patient has already been started on Xgeva. For other medically accepted indication, no trial of alternatives is required.

XIFAXAN

Products Affected

- Xifaxan TABS 550MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

XIAFLEX

Products Affected

- Xiaflex

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Xiaflex is authorized for the treatment of a symptomatic Dupuytren's contracture in adults when there is both a palpable cord and a functional impairment as manifested by a metacarpophalangeal (MCP) joint or proximal interphalangeal (PIP) joint contracture of 20 degrees or greater.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	B vs D coverage determination

XYREM

Products Affected

- Xyrem

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	The patient must not be taking any sedative hypnotic agents or other CNS depressants.
Required Medical Information	Documentation of diagnosis, sleep study
Age Restrictions	7 years of age and older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	For the initial treatment of excessive daytime sleepiness (EDS) in patients with narcolepsy: trial and failure, contraindication, or intolerance to one CNS stimulant (e.g., methylphenidate, dexamethylphenidate, dextroamphetamine) or armodafinil is required. For continuation of therapy in the treatment of excessive daytime sleepiness (EDS) in patients with narcolepsy, approve if patient has already been started on Xyrem. For other medically accepted indication, no trial of alternatives is required.

ZTLIDO

Products Affected

- Ztlido

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Diabetic neuropathic pain, cancer related neuropathy
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

PART B VERSUS PART D

Products Affected

- Acetylcysteine SOLN
- Acyclovir Sodium INJ 50MG/ML
- Albuterol Sulfate NEBU
- Aminosyn INJ 148MEQ/L;
1280MG/100ML; 980MG/100ML;
1280MG/100ML; 300MG/100ML;
720MG/100ML; 940MG/100ML;
720MG/100ML; 400MG/100ML;
440MG/100ML; 860MG/100ML;
420MG/100ML; 520MG/100ML;
160MG/100ML; 44MG/100ML;
800MG/100ML, 90MEQ/L;
1100MG/100ML; 850MG/100ML;
35MEQ/L; 1100MG/100ML;
260MG/100ML; 620MG/100ML;
810MG/100ML; 624MG/100ML;
340MG/100ML; 380MG/100ML;
750MG/100ML; 370MG/100ML;
460MG/100ML; 150MG/100ML;
44MG/100ML; 680MG/100ML
- Aminosyn II INJ 107.6MEQ/L;
1490MG/100ML; 1527MG/100ML;
1050MG/100ML; 1107MG/100ML;
750MG/100ML; 450MG/100ML;
990MG/100ML; 1500MG/100ML;
1575MG/100ML; 258MG/100ML;
447MG/100ML; 1083MG/100ML;
795MG/100ML; 50MEQ/L;
600MG/100ML; 300MG/100ML;
405MG/100ML; 750MG/100ML,
71.8MEQ/L; 993MG/100ML;
1018MG/100ML; 700MG/100ML;
738MG/100ML; 500MG/100ML;
300MG/100ML; 660MG/100ML;
1000MG/100ML; 1050MG/100ML;
172MG/100ML; 298MG/100ML;
722MG/100ML; 530MG/100ML;
38MEQ/L; 400MG/100ML;
200MG/100ML; 270MG/100ML;
500MG/100ML

- Aminosyn M INJ 65MEQ/L;
448MG/100ML; 343MG/100ML;
40MEQ/L; 448MG/100ML;
105MG/100ML; 252MG/100ML;
329MG/100ML; 252MG/100ML;
3MEQ/L; 140MG/100ML;
154MG/100ML; 3.5MMOLE/L;
13MEQ/L; 300MG/100ML;
147MG/100ML; 40MEQ/L;
182MG/100ML; 56MG/100ML;
31MG/100ML; 280MG/100ML
- Aminosyn-pf INJ 46MEQ/L;
698MG/100ML; 1227MG/100ML;
527MG/100ML; 820MG/100ML;
385MG/100ML; 312MG/100ML;
760MG/100ML; 1200MG/100ML;
677MG/100ML; 180MG/100ML;
427MG/100ML; 812MG/100ML;
495MG/100ML; 3.4MEQ/L;
70MG/100ML; 512MG/100ML;
180MG/100ML; 44MG/100ML;
673MG/100ML
- Aminosyn-pf 7%
- Aprepitant
- Arsenic Trioxide INJ
- Bendeka
- Budesonide SUSP
- Busulfan
- Busulfex
- Cerezyme
- Clinimix 4.25%/dextrose 10%
- Clinimix 4.25%/dextrose 25%
- Clinimix 4.25%/dextrose 5%
- Clinimix 5%/dextrose 15%
- Clinimix 5%/dextrose 20%
- Clinimix 5%/dextrose 25%

- Clinimix E 2.75%/dextrose 10% INJ
570MG/100ML; 317MG/100ML;
33MG/100ML; 10GM/100ML;
283MG/100ML; 132MG/100ML;
165MG/100ML; 201MG/100ML;
159MG/100ML; 51MG/100ML;
110MG/100ML; 454MG/100ML;
154MG/100ML; 261MG/100ML;
187MG/100ML; 138MG/100ML;
217MG/100ML; 112MG/100ML;
116MG/100ML; 50MG/100ML;
11MG/100ML; 160MG/100ML
- Clinimix E 4.25%/dextrose 10%
- Clinimix E 4.25%/dextrose 25%
- Clinimix N9g15e
- Clinisol Sf 15%
- Cromolyn Sodium NEBU
- Cyclophosphamide CAPS
- Cyclophosphamide INJ
- Daptomycin
- Daunorubicin Hydrochloride
- Dextrose INJ 50%
- Dextrose 10%/nacl 0.45%
- Dextrose 5% /electrolyte #48 Viaflex
- Dextrose 10%
- Dextrose 10%/nacl 0.2%
- Dextrose 2.5%/nacl 0.45%
- Dextrose 20%
- Dextrose 25% INJ 250MG/ML
- Dextrose 30%
- Dextrose 40%
- Dextrose 5%/lactated Ringers
- Dextrose 50%
- Duramorph
- Emend SUSR
- Engerix-b
- Etoposide INJ 100MG/5ML,
1GM/50ML, 500MG/25ML
- Fabrazyme
- Faslodex INJ 250MG/5ML
- Firmagon
- Fludarabine Phosphate
- Folutyn
- Freamine Hbc 6.9%

- Freamine III INJ 89MEQ/L;
710MG/100ML; 950MG/100ML;
3MEQ/L; 24MG/100ML;
1400MG/100ML; 280MG/100ML;
690MG/100ML; 910MG/100ML;
730MG/100ML; 530MG/100ML;
560MG/100ML; 10MMOLE/L;
120MG/100ML; 1120MG/100ML;
590MG/100ML; 10MEQ/L;
400MG/100ML; 150MG/100ML;
660MG/100ML
- Fulvestrant
- Gamunex-c
- Gemcitabine
- Gemcitabine Hcl
- Gemcitabine Hydrochloride INJ
1GM, 1GM/26.3ML, 200MG/2ML,
200MG/5.26ML, 2GM/20ML,
2GM/52.6ML
- Granisetron Hcl TABS
- Hepatamine
- Hizentra
- Humulin R U-500 (concentrated)
- Imovax Rabies (h.d.c.v.)
- Infugem
- Intralipid INJ 20GM/100ML,
30GM/100ML
- Ipratropium Bromide INHALATION
SOLN 0.02%
- Ipratropium Bromide/albuterol
Sulfate
- Irinotecan INJ 100MG/5ML,
500MG/25ML
- Irinotecan Hcl
- Irinotecan Hydrochloride
- Kabiven
- Kcl 0.075%/d5w/nacl 0.45%
- Kcl 0.15%/d5w/nacl 0.2%
- Kcl 0.15%/d5w/nacl 0.225% INJ 5%;
20MEQ/L; 0.225%
- Kcl 0.15%/d5w/nacl 0.45%
- Kcl 0.15%/d5w/nacl 0.9%
- Kcl 0.3%/d5w/nacl 0.45%
- Kcl 0.3%/d5w/nacl 0.9%
- Kyprolis

- Lactated Ringers INJ 3MEQ/L;
109MEQ/L; 28MEQ/L; 4MEQ/L;
130MEQ/L
- Lactated Ringers Viaflex
- Magnesium Sulfate INJ
20GM/500ML, 2GM/50ML,
40GM/1000ML, 4GM/100ML,
4GM/50ML, 50%
- Magnesium Sulfate In D5w INJ 5%;
1GM/100ML
- Melphalan Hydrochloride
- Mesna
- Morphine Sulfate INJ 0.5MG/ML,
10MG/ML, 150MG/30ML,
1MG/ML, 2MG/ML, 4MG/ML,
5MG/ML, 8MG/ML
- Nebupent
- Nephramine
- Normosol -r
- Normosol-m In D5w
- Normosol-r
- Normosol-r In D5w
- Nutrilipid
- Ondansetron Hcl SOLN
- Ondansetron Hcl TABS 24MG
- Ondansetron Hydrochloride TABS
- Ondansetron Odt
- Paclitaxel INJ 100MG/16.7ML,
150MG/25ML, 300MG/50ML,
30MG/5ML
- Pamidronate Disodium
- Perforomist
- Perikabiven
- Plenamine
- Potassium Chloride INJ
10MEQ/100ML, 20MEQ/100ML,
2MEQ/ML, 40MEQ/100ML
- Potassium Chloride/dextrose INJ 5%;
20MEQ/L, 5%; 40MEQ/L
- Potassium Chloride/dextrose/lactated
Ringers
- Potassium Chloride/dextrose/sodium
Chloride

- Potassium Chloride/sodium Chloride
INJ 20MEQ/L; 0.45%, 20MEQ/L;
0.9%, 40MEQ/L; 0.9%
- Prednisone TABS 10MG, 1MG,
2.5MG, 20MG, 50MG, 5MG
- Premasol
- Procalamine
- Prolastin-c
- Prosol
- Pulmozyme
- Rabavert
- Recombivax Hb
- Remodulin
- Ringers Injection INJ 4.5MEQ/L;
156MEQ/L; 4MEQ/L; 147MEQ/L
- Simulect
- Sivextro INJ
- Sodium Lactate INJ 5MEQ/ML
- Temsirolimus
- Tobramycin NEBU
- Toposar INJ 100MG/5ML,
1GM/50ML, 500MG/25ML
- Torisel
- Tpn Electrolytes
- Travasol INJ 52MEQ/L;
1760MG/100ML; 880MG/100ML;
34MEQ/L; 1760MG/100ML;
372MG/100ML; 406MG/100ML;
526MG/100ML; 492MG/100ML;
492MG/100ML; 526MG/100ML;
356MG/100ML; 500MG/100ML;
356MG/100ML; 390MG/100ML;
34MG/100ML; 152MG/100ML
- Treanda INJ 100MG, 25MG
- Treprostinil
- Trisenox INJ 12MG/6ML
- Trogarzo
- Trophamine
- Vincasar Pfs
- Vincristine Sulfate INJ
- Vinorelbine Tartrate
- Vyxeos
- Zoledronic Acid INJ 4MG/5ML,
5MG/100ML

Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

INDEX

A

Abelcet	7
Abiraterone Acetate.....	94
Abraxane	10
Acetylcysteine	152
Acitretin.....	1
Actimmune	2
Acyclovir Sodium	152
Adagen	70
Adempas.....	138
Afinitor.....	94
Afinitor Disperz.....	94
Albuterol Sulfate	152
Aldurazyme	40
Alecensa	94
Alimta.....	3
Aliqopa	10
Alosetron	4
Alosetron Hydrochloride.....	4
Alunbrig	94
Ambisome	7
Ambrisentan	138
Aminosyn	152
Aminosyn II.....	152
Aminosyn M.....	153
Aminosyn-pf.....	153
Aminosyn-pf 7%	153
Amitriptyline Hcl	65
Amitriptyline Hydrochloride.....	65
Amphotericin B.....	7
Anabolic Steroids, Androgens.....	5
Anadrol-50	5
Antifungals, Azole	6
Antifungals, Polyene	7
Antifungals, Superficial And Systemic.....	8
Antifungals, Triazole.....	9
Antineoplastics, Monoclonal Antibodies	10

Apokyn.....	12
Aprepitant	153
Aranesp Albumin Free.....	51
Arcalyst.....	13
Arikayce.....	14
Armodafinil.....	101
Arsenic Trioxide	153
Astagraf XL	77
Auryxia	15
Austedo	16
Avastin	10
Avita.....	31
Avonex	79
Avonex Pen.....	79
Azathioprine.....	77

B

Balversa.....	94
Banzel	17
Bavencio	10
Bendeka.....	153
Benlysta.....	18
Benzotropine Mesylate	55
Besponsa	10
Betaseron.....	79
Bexarotene	19
Bortezomib.....	10
Bosulif.....	94
Botox.....	20
Braftovi	94
Brukina	21
Budesonide.....	153
Buprenorphine.....	22
Buprenorphine Hcl.....	22
Busulfan	153
Busulfex	153
Butalbital/acetaminophen/caffeine	56

C	
Cabometyx	94
Calquence	94
Caprelsa.....	94
Carbaglu	23
Casprofungin Acetate	8
Cayston.....	24
Cerezyme.....	153
Chorionic Gonadotropin.....	53
Cinryze	25
Clinimix 4.25%/dextrose 10%	153
Clinimix 4.25%/dextrose 25%	153
Clinimix 4.25%/dextrose 5%	153
Clinimix 5%/dextrose 15%	153
Clinimix 5%/dextrose 20%	153
Clinimix 5%/dextrose 25%	153
Clinimix E 2.75%/dextrose 10%.....	154
Clinimix E 4.25%/dextrose 10%.....	154
Clinimix E 4.25%/dextrose 25%.....	154
Clinimix N9g15e.....	154
Clinisol Sf 15%	154
Clomipramine Hcl.....	65
Colony Stimulating Factors.....	26
Cometriq.....	94
Copaxone.....	79
Copiktra.....	94
Corlanor.....	27
Cotellic	94
Cromolyn Sodium	154
Cyclobenzaprine Hydrochloride	64
Cyclophosphamide	154
Cyclosporine.....	77
Cyclosporine Modified.....	77
Cyramza	10
Cystaran.....	28

D

Dalfampridine.....	29
Dalfampridine Er.....	29
Daliresp	112
Daptomycin	154
Darzalex	10
Daunorubicin Hydrochloride.....	154
Daurismo	94

Demser	30
Dermatological Retinoids	31
Dermatological Wound Care Agents	32
Dextrose	154
Dextrose 10%/nacl 0.45%.....	154
Dextrose 5% /electrolyte #48 Viaflex.....	154
Dextrose 10%.....	154
Dextrose 10%/nacl 0.2%.....	154
Dextrose 2.5%/nacl 0.45%.....	154
Dextrose 20%.....	154
Dextrose 25%.....	154
Dextrose 30%.....	154
Dextrose 40%.....	154
Dextrose 5%/lactated Ringers.....	154
Dextrose 50%.....	154
Dihydroergotamine Mesylate.....	33
Dipyridamole	61
Dotti	57
Doxepin Hcl.....	65
Doxepin Hydrochloride	65
Dronabinol	34
Duavee	35
Dupixent.....	36
Dupixent.....	36
Duramorph	154

E

Elaprase.....	40
Emend	154
Enbrel.....	71
Enbrel Mini	71
Enbrel Sureclick.....	71
Endocrine And Metabolic Agents.....	39
Engerix-b.....	154
Enhertu.....	10
Enzyme Replacement/modifiers	40
Epclusa	41
Epidiolex	42
Erivedge	43
Erleada	44
Erlotinib Hydrochloride	94
Esbriet	69
Esgic.....	56
Estradiol	57
Etoposide.....	154

Everolimus (antineoplastic).....	94
Evomela.....	45
Eylea.....	46

F

Fabrazyme	154
Farydak.....	94
Faslodex	154
Fentanyl Citrate Oral Transmucosal	134
Firmagon	154
Fludarabine Phosphate	154
Folotyn	154
Forteo	92
Freamine Hbc 6.9%.....	154
Freamine III.....	155
Fulvestrant.....	155
Fyavolv.....	57
Fycompa.....	47

G

Gamunex-c	155
Gattex	48
Gazyva.....	10
Gemcitabine	155
Gemcitabine Hcl.....	155
Gemcitabine Hydrochloride	155
Gengraf.....	77
Genotropin.....	113
Genotropin Miniquick.....	113
Gilenya	79
Gilotrif.....	94
Granisetron Hcl	155

H

Halaven.....	49
Harvoni.....	50
Hematopoietics.....	51
Hepatamine.....	155
Herceptin	10
Herceptin Hylecta.....	10
Hetlioz	52
Hizentra	155
Hormonal Agents, Gonadotropins.....	53
Hormonal Agents, Somatostatin Analogs	54
Hrm - Benzotropine.....	55

Hrm - Butalbital Combinations.....	56
Hrm - Estrogens	57
HRM - first generation antihistamines.....	58
Hrm - Megestrol.....	59
Hrm - Perphenazine/amitriptyline.....	60
Hrm - Platelet Modifying Agents	61
Hrm - Promethazine.....	62
Hrm - Skeletal Muscle Relaxants	64
Hrm - Tricyclic Antidepressants.....	65
Humira	71
Humira Pediatric Crohns Disease Starter Pack....	71
Humira Pen	71
Humira Pen-cd/uc/hs Starter	71
Humira Pen-ps/uv Starter.....	71
Humulin R U-500 (concentrated)	155
Hydroxyprogesterone.....	67
Hydroxyprogesterone Caproate	67
Hydroxyzine Hcl.....	58

I

Ibrance.....	94
Icatibant.....	68
Icatibant Acetate	68
Iclusig.....	94
Idhifa	94
Idiopathic Pulmonary Fibrosis.....	69
Imatinib Mesylate	94
Imbruvica	94
Imfinzi.....	10
Imipramine Hcl	65
Imipramine Hydrochloride.....	65
Immune Stimulants	70
Immune Suppressants	71
Immune Suppressants - Rinvoq	74
Immune Suppressants - Stelara.....	75
Immune Suppressants - Transplant Related.....	77
Immune Suppressants - Xeljanz.....	78
Immunomodulators	79
Imovax Rabies (h.d.c.v.).....	155
Increlex	83
Infliximab.....	80
Infugem	155
Inlyta	94
Inrebic	94
Insulin-like Growth Factor.....	83

Intralipid.....	155
Ipratropium Bromide.....	155
Ipratropium Bromide/albuterol Sulfate.....	155
Iressa.....	94
Irinotecan.....	155
Irinotecan Hcl.....	155
Irinotecan Hydrochloride	155
Istodax	84
Istodax (overfill).....	84
Itraconazole	8

J

Jadenu.....	85
Jakafi	94

K

Kabiven	155
Kadcyla.....	10
Kalydeco.....	86
Kanjinti.....	10
Kcl 0.075%/d5w/nacl 0.45%.....	155
Kcl 0.15%/d5w/nacl 0.2%.....	155
Kcl 0.15%/d5w/nacl 0.225%.....	155
Kcl 0.15%/d5w/nacl 0.45%.....	155
Kcl 0.15%/d5w/nacl 0.9%.....	155
Kcl 0.3%/d5w/nacl 0.45%.....	155
Kcl 0.3%/d5w/nacl 0.9%.....	155
Keytruda.....	10
Kisqali	94
Kisqali Femara 200 Dose	94
Kisqali Femara 400 Dose	94
Kisqali Femara 600 Dose	94
Korlym	87
Kuvan	88
Kyprolis.....	155

L

Lactated Ringers.....	156
Lactated Ringers Viaflex.....	156
Lartruvo.....	10
Lenvima 10 Mg Daily Dose.....	94
Lenvima 12mg Daily Dose	94
Lenvima 14 Mg Daily Dose.....	94
Lenvima 18 Mg Daily Dose.....	94
Lenvima 20 Mg Daily Dose.....	94

Lenvima 24 Mg Daily Dose.....	94
Lenvima 4 Mg Daily Dose.....	94
Lenvima 8 Mg Daily Dose.....	94
Letairis	138
Leuprolide Acetate.....	113
Libtayo	10
Lidocaine.....	89
Lidocaine Patch.....	89
Lonsurf.....	94
Lorbrena.....	94
Lumizyme	90
Lumoxiti.....	10
Lupron Depot (1-month).....	113
Lupron Depot (3-month).....	113
Lupron Depot (4-month).....	113
Lupron Depot (6-month).....	113
Lupron Depot-ped (1-month).....	113
Lupron Depot-ped (3-month).....	113
Lynparza	95

M

Magnesium Sulfate	156
Magnesium Sulfate In D5w	156
Mavyret.....	91
Megestrol Acetate	59
Mekinist	95
Mektovi.....	95
Melphalan Hydrochloride	156
Memantine Hcl Titration Pak.....	100
Memantine Hydrochloride	100
Memantine Hydrochloride Er	100
Mesna.....	156
Metabolic Bone Disease Agents.....	92
Methocarbamol	64
Molecular Target Inhibitors	93
Monoclonal Antibodies.....	97
Morphine Sulfate	156
Mvasi.....	10
Mycophenolate Mofetil.....	77
Mycophenolic Acid Dr	77
Mylotarg.....	10

N

Naglazyme	40
Namzaric.....	100

Natpara	98	Perforomist.....	156
Nayzilam	99	Perikabiven	156
Nebupent	156	Perjeta	10
Nephramine	156	Perphenazine/amitriptyline	60
Nerlynx.....	95	Phosphodiesterase Type 4 (pde4) Inhibitors.....	112
Neulasta.....	26	Piqray 200mg Daily Dose.....	95
Neulasta Onpro.....	26	Piqray 250mg Daily Dose.....	95
Nexavar	95	Piqray 300mg Daily Dose.....	95
Ninlaro.....	95	Pituitary Hormones	113
Nmda Receptor Antagonist.....	100	Plenaminate.....	156
Non-amphetamine Central Nervous System Agents	101	Pomalyst.....	95
.....	101	Posaconazole Dr.....	9
Normosol -r	156	Potassium Chloride	156
Normosol-m In D5w	156	Potassium Chloride/dextrose	156
Normosol-r	156	Potassium Chloride/dextrose/lactated Ringers ..	156
Normosol-r In D5w	156	Potassium Chloride/dextrose/sodium Chloride..	156
Northera.....	102	Potassium Chloride/sodium Chloride	157
Noxafil.....	9	Poteligeo	10
Nubeqa	113	Prednisone.....	157
Nuedexta.....	103	Premarin.....	57
Nulojix.....	104	Premasol.....	157
Nuplazid	105	Procalamine.....	157
Nutrilipid.....	156	Prograf.....	77
O		Prolastin-c	157
Ocaliva	106	Promacta	114
Ocrevus.....	107	Promethazine Hcl.....	62
Octreotide Acetate.....	54	Promethazine Hcl Plain.....	62
Odomzo	108	Promethazine Hydrochloride	62
Ofev.....	69	Prosol	157
Ondansetron Hcl.....	156	Pulmozyme	157
Ondansetron Hydrochloride	156	Purixan	115
Ondansetron Odt	156	Q	
Opdivo.....	10	Quinine Sulfate	116
Opsumit.....	138	Quninie Sulfate	116
Orencia	109	R	
Orencia Clickject.....	109	Rabavert	157
Orkambi.....	110	Rapamune	77
Oxandrolone.....	5	Recombivax Hb	157
P		Regranex	32
Paclitaxel.....	156	Relistor.....	117
Padcev	111	Remodulin.....	157
Pamidronate Disodium.....	156	Renflexis	80
Part B Versus Part D	151	Repatha	118

Repatha Pushtronex System.....	118
Repatha Sureclick.....	118
Retacrit.....	51
Revlimid.....	120
Ringers Injection.....	157
Rinvoq.....	74
Rituxan.....	10
Rituxan Hycela.....	10
Romidepsin.....	84
Rozlytrek.....	121
Rubraca.....	95
Ruconest.....	122
Rydapt.....	95

S

Samsca.....	123
Sandimmune.....	77
Signifor.....	124
Sildenafil Citrate.....	138
Simulect.....	157
Sirolimus.....	77
Sirturo.....	125
Sivextro.....	157
Skyrizi.....	71
Sodium Lactate.....	157
Sodium Phenylbutyrate.....	126
Somatuline Depot.....	54
Somavert.....	39
Sprycel.....	95
Stelara.....	75
Stivarga.....	95
Sutent.....	95
Sylatron.....	127
Sympazan.....	128
Synagis.....	129
Synarel.....	113
Synribo.....	95

T

Tacrolimus.....	77
Tafinlar.....	95
Tagrisso.....	95
Talzenna.....	95
Targretin.....	130
Targretin (topical).....	130

Tasigna.....	95
Tecentriq.....	10
Tecfidera.....	79
Tecfidera Starter Pack.....	79
Temsirolimus.....	157
Testosterone.....	5
Testosterone Pump.....	5
Tetrabenazine.....	131
Thalidomide (thalomid).....	132
Thalomid.....	132
Thiotepa.....	133
Tibsovo.....	95
Tobramycin.....	157
Toposar.....	157
Torisel.....	157
Tpn Electrolytes.....	157
Transmucosal Fentanyl Citrate.....	134
Travasol.....	157
Treanda.....	157
Trelstar Mixject.....	113
Treprostinil.....	157
Tretinoin.....	31
Tretinoin Microsphere.....	31
Tretinoin Microsphere Pump.....	31
Trimipramine Maleate.....	65
Triptodur.....	113
Trisenox.....	157
Trogarzo.....	157
Trophamine.....	157
Tykerb.....	95
Tymlos.....	92
Tysabri.....	135

U

Unituxin.....	10
Uptravi.....	136

V

Valchlor.....	137
Valchlor Gel.....	137
Vasodilators.....	138
Vectibix.....	10
Velcade.....	10
Venclexta.....	95
Venclexta Starting Pack.....	95

Ventavis.....	139
Verzenio	95
Viberzi.....	140
Vigabatrin.....	141
Vigadrone.....	141
Vincasar Pfs.....	157
Vincristine Sulfate.....	157
Vinorelbine Tartrate	157
Vitrakvi.....	142
Vivitrol	143
Vizimpro.....	95
Voriconazole	6
Vosevi.....	144
Votrient.....	95
Vyxeos.....	157

X

Xalkori.....	95
Xatmep	145
Xeljanz	78
Xeljanz Xr	78
Xgeva	146
Xiaflex.....	148
Xifaxan.....	147

Xolair	97
Xospata	95
Xpovio 100 Mg Once Weekly	95
Xpovio 60 Mg Once Weekly	95
Xpovio 80 Mg Once Weekly	95
Xpovio 80 Mg Twice Weekly.....	95
Xtandi.....	95
Xyrem	149

Y

Yervoy.....	10
Yondelis	10

Z

Zarxio	26
Zebutal	56
Zejula	95
Zelboraf.....	95
Zoledronic Acid	157
Zortress	77
Ztlido.....	150
Zydelig	95
Zykadia	95
Zytiga	95